

EVALUATION OF ANTI-GOUT POTENTIAL OF SELECTED MALAYSIAN  
MEDICINAL PLANTS

FAZLEEN IZZANY BINTI ABU BAKAR

A thesis submitted in  
fulfillment of the requirement for the award of the  
Doctor of Philosophy in Science

Faculty of Applied Sciences and Technology  
Universiti Tun Hussein Onn Malaysia

JULY 2020

## DEDICATION

Dedicated to: My late father, Abu Bakar Talib, my mother, Suriyah Mohd Isa, My siblings, Ahmad Fikri, Fariza Ilyani, Ahmad Faizal Afiffy (late brother) and Fazreena Aryanni, my lecturers and friends for their support and encouragement over the years who understand and guided me through the journey of my study and for all who believe in me and keep whispering “You can do it”.



PTTA UTHM  
PERPUSTAKAAN TUNKU TUN AMINAH

## ACKNOWLEDGEMENT

In the name of Allah, The Most Gracious and The Most Merciful

I would like to express the sincerest gratitude to my supervisor, Assoc. Prof. Ts. Dr. Mohd Fadzelly Abu Bakar, co supervisors, Ts. Dr. Norazlin Abdullah and Prof. Dr. Susi Endrini for the continuous support and motivation provided throughout my studies. Their guidance and generosity in providing help and knowledge throughout this journey has amazingly inspired me.

I would also like to thank my family: Mak, Along, Angah and Ina for their support, trust, love, prayer, and financial aid. Sacrifices were made to ensure my success, and I am forever grateful. Apart from that, special thanks to all the laboratory assistants of Department of Technology and Natural Resources, Faculty of Applied Sciences and Technology, Universiti Tun Hussein Onn Malaysia (UTHM) especially Mr. Ishak Bin Ayub. I would also like to extend my deepest appreciation to my best friend, Nurulain Syuhada Mohd Yazid for helping me throughout my PhD journey. Thank you for the love and friendship. Biggest thanks to all the research assistants in Universitas Yarsi, Jakarta, Indonesia especially Silviatun Nihayah, Indah Permata Yuda and Arif Ashari for their help.

Finally, this research would not be complete without the financial support and I am grateful to the Ministry of Education for the Fundamental Research Grant Scheme (FRGS Vot K099), Universiti Tun Hussein Onn Malaysia for Postgraduate Research Grant (GPPS Vot U758) and Yayasan Negeri Sembilan that immensely helped in funding this study.

To everyone else who have directly and indirectly contributed in the success of this study, I humbly thank all of you. Only Allah SWT will be able to repay your kindness.

## ABSTRACT

Gout is a type of arthritis that causes painful inflammation in joints due to the elevation of uric acid in the blood. Allopurinol is a common drug used for treating gout. However, due to unwanted side effects, new alternatives with fewer side effects are desired. The objectives of the study are to screen the six potential Malaysian medicinal plants for anti-gout activity using *in vitro* enzyme assay; to optimize the extraction conditions of the three selected plants using response surface methodology (RSM); to determine the cell viability of the three extracts towards RAW 264.7 murine macrophage cell line; to investigate the *in vivo* anti-gout mechanism of three selected plants; and to profile the bioactive compounds present in the three selected plant extracts. Six selected plants were screened for their anti-gout activity *in vitro* of which three plants (i.e *Euphorbia hirta*, *Strobilanthes crispus* and *Christia vespertilionis*) were further analyzed for their optimum extraction conditions using RSM. The effects of the plant extracts on the serum uric acid levels in hyperuricemic rats were then evaluated and lastly the bioactive compounds of the plant extracts were analyzed using liquid chromatography-mass spectrometry (LC-MS). *E. hirta*, *S. crispus* and *C. vespertilionis* exhibited XO inhibitory activity *in vitro* with more than 60 % at 100 µg/mL. Findings showed that the optimum extraction conditions of phytochemical compounds and anti-gout activity of these plants were at temperature of 65.09-79.07 °C for 5.0-17.42 min with solid to liquid ratio of 1:17.33-1:20 g/mL, respectively. *C. vespertilionis* displayed less cytotoxic against RAW264.7 cells as compared to *S. crispus* and *E. hirta*. For the *in vivo* study, the treatment of the hyperuricemic rats with 200 mg/kg of *E. hirta*, *S. crispus* and *C. vespertilionis* extracts reduced the serum uric acid levels significantly by more than 40 % as compared to hyperuricemic rats. The LC-MS analysis also revealed the presence of phytochemicals such as flavonoid and phenolic compounds in the plant extracts which might be associated with the anti-gout activity. Hence, this study clearly demonstrated that *E. hirta*, *S. crispus* and *C. vespertilionis* aqueous extracts are potential as anti-gout agents.



## ABSTRAK

Gout adalah sejenis arthritis yang menyebabkan keradangan yang menyakitkan di sendi disebabkan oleh peningkatan asid urik dalam darah. Allopurinol ialah ubat yang biasa digunakan untuk merawat gout. Walau bagaimanapun, disebabkan kesan sampingan yang tidak diingini, alternatif baru dengan kesan sampingan yang lebih sedikit adalah diperlukan. Objektif kajian ini adalah untuk mengenal pasti enam tumbuhan ubat-ubatan Malaysia yang berpotensi sebagai anti-gout dengan menggunakan enzim *in vitro*; untuk mengoptimumkan keadaan ekstraksi tiga tumbuhan terpilih dengan menggunakan kaedah permukaan respon (KSR); untuk menentukan daya maju sel tiga ekstrak terhadap sel makrofaj RAW 264.7; untuk mengkaji mekanisme aktiviti anti-gout *in vivo* oleh tiga tumbuhan terpilih; dan membuat profiling sebatian bioaktif yang terdapat di dalam tiga ekstrak tumbuhan terpilih. Enam tumbuhan yang dipilih telah dikaji untuk aktiviti anti-gout di mana tiga tumbuhan (i.e *Euphorbia hirta*, *Strobilanthes crispus* dan *Christia vespertilionis*) dianalisa lagi untuk keadaan pengekstrakan optimum menggunakan KSR. Kesan ekstrak tumbuhan pada paras asid urik dalam darah tikus kemudian dinilai dan akhirnya sebatian bioaktif ekstrak tumbuhan dianalisa menggunakan spektrometri jisim-kromatografi cecair (SJ-KC). *E. hirta*, *S. crispus* dan *C. vespertilionis* menunjukkan aktiviti perencatan XO secara *in vitro* pada kadar lebih 60 % pada kepekatan 100 µg/mL. Keputusan menunjukkan bahawa keadaan pengekstrakan optimum bagi tumbuhan ini adalah pada suhu 65.09-79.07 °C untuk 5.0-17.4 minit dengan nisbah pepejal kepada cecair 1:17.33-1:20 g/mL, masing-masing. *C. vespertilionis* menunjukkan ketoksikan yang rendah terhadap sel RAW264.7 berbanding dengan *S. crispus* dan *E. hirta*. Untuk kajian terhadap haiwan, tikus yang diberikan 200 mg/kg ekstrak *E. hirta*, *S. crispus* dan *C. vespertilionis* mengurangkan paras serum asid urik secara signifikan sebanyak lebih daripada 40 % berbanding tikus kawalan. Analisis SJ-KC juga mendedahkan kehadiran fitokimia seperti flavonoid dan fenolik dalam ekstrak tumbuhan yang mungkin boleh dikaitkan dengan aktiviti anti-

gout. Secara keseluruhan, kajian ini jelas menunjukkan ekstrak akueus *E. hirta*, *S. crista* dan *C. vespertilionis* sebagai agen anti-gout yang berpotensi.



## TABLE OF CONTENTS

<b>TITLE</b>	<b>i</b>
<b>DECLARATION</b>	<b>ii</b>
<b>DEDICATION</b>	<b>iii</b>
<b>ACKNOWLEDGEMENT</b>	<b>iv</b>
<b>ABSTRACT</b>	<b>v</b>
<b>ABSTRAK</b>	<b>vi</b>
<b>TABLE OF CONTENTS</b>	<b>viii</b>
<b>LIST OF TABLES</b>	<b>xiv</b>
<b>LIST OF FIGURES</b>	<b>xvii</b>
<b>LIST OF SYMBOLS AND ABBREVIATIONS</b>	<b>xxi</b>
<b>LIST OF APPENDICES</b>	<b>xxiv</b>
<b>LIST OF PUBLICATIONS</b>	<b>xxvi</b>
<b>LIST OF AWARDS</b>	<b>xxviii</b>
<b>CHAPTER 1 GENERAL INTRODUCTION</b>	<b>1</b>
1.1 Introduction	1
1.2 Background of the study	1
1.3 Problem statement	4
1.4 Objectives of the study	5
1.5 Significance of research	6
1.6 Scope and limitations of study	7
<b>CHAPTER 2 LITERATURE REVIEW</b>	<b>8</b>
2.1 Gout	8

2.1.1	Epidemiology of gout	11
2.1.2	Factors of gout	12
2.1.2.1	Age	12
2.1.2.2	Gender	12
2.1.2.3	Diet and alcohol intake	13
2.1.2.4	Obesity	14
2.1.2.5	Genetics	14
2.1.3	Current drugs and treatment for gout	15
2.1.3.1	Nonsteroidal anti-inflammatory drugs (NSAIDs)	15
2.1.3.2	Colchicine	15
2.1.3.3	Corticosteroids	17
2.1.3.4	Xanthine oxidase inhibitors	17
2.1.3.5	Uricosuric agents	19
2.1.4	Pathogenesis of gout	20
2.1.5	Uric acid	21
2.1.6	Mechanisms of drugs in inhibiting xanthine oxidase (XO) activity	23
2.2	Potential Malaysian medicinal plants as alternative to treat gout	24
2.2.1	<i>Moringa oleifera</i>	25
2.2.2	<i>Strobilanthes crispus</i>	26
2.2.3	<i>Artocarpus altilis</i>	27
2.2.4	<i>Ortosiphon aristatus</i>	28
2.2.5	<i>Euphorbia hirta</i>	29
2.2.6	<i>Christia vespertilionis</i>	30

2.3	Malaysian medicinal plants for anti-gout activity in <i>in vitro</i> study	31
2.3.1	Enzyme assay	31
2.3.2	Cell study	33
2.4	Malaysian medicinal plants for anti-gout activity in <i>in vivo</i> study	42
2.5	Compounds that contribute to anti-gout effects	51
2.6	Response surface methodology	51
2.6.1	Factorial and central composite design	53
2.6.2	Box-behnken design	54
2.6.3	Application of RSM in optimizing the extraction conditions of plants for maximum phytochemicals yields	55
2.7	Summary	56
<b>CHAPTER 3 METHODOLOGY</b>		<b>57</b>
3.1	Study design	57
3.2	Sample preparation	57
3.3	Sample extraction	59
3.4	Screening of anti-gout potential of selected Malaysian medicinal plants <i>in vitro</i> and phytochemical content determination	59
3.4.1	Xanthine oxidase inhibitory activity assay	59
3.4.2	IC <sub>50</sub> concentration determination	60
3.4.3	Determination of total phenolic content	60
3.4.4	Determination of total flavonoid content	60
3.5	Optimization of extraction conditions using response surface methodology (RSM)	61

3.5.1	Screening of important variables	61
3.5.2	Experimental design	61
3.5.3	Model validation	62
3.6	Cell study	63
3.6.1	Cell culture and treatment	63
3.6.2	Cell viability determination	63
3.7	<i>In vivo</i> anti-gout study	64
3.7.1	Animals	64
3.7.2	Oral acute toxicity study	64
3.7.3	Test compound preparation	65
3.7.4	Induction of hyperuricemia	65
3.7.5	Experimental design and treatment of animals	65
3.7.6	Liver function test	67
3.7.7	Preparation of liver homogenate	67
3.7.8	Assay of XO activity in rat liver	68
3.8	Phytochemical profiling of selected crude extracts	68
3.8.1	Liquid Chromatography-Mass Spectrometry (LC-MS) analysis	68
3.9	Statistical analysis	69
<b>CHAPTER 4 RESULTS AND DISCUSSION</b>		<b>70</b>
4.1	Screening of xanthine oxidase (XO) inhibitory activity of selected Malaysian medicinal plants	70
4.1.1	Total phenolic content (TPC)	74
4.1.2	Total flavonoid content (TFC)	77

4.1.3	Correlation between total phenolic content (TPC), total flavonoid content (TFC) and XO inhibitory activity	80
4.2	Optimization of plant extract using response surface methodology (RSM)	81
4.2.1	Screening the important process variables using a pareto chart	81
4.2.2	Fitting for the best models of <i>E. hirta</i> , <i>S. crispa</i> and <i>C. vespertilionis</i>	86
4.2.3	Modelling of total flavonoid content of <i>E. hirta</i> , <i>S. crispa</i> and <i>C. vespertilionis</i>	89
4.2.4	Combined effect of extraction time, temperature and solid to liquid ratio on total flavonoid content	92
4.2.5	Modelling of total phenolic content of <i>E. hirta</i> , <i>S. crispa</i> and <i>C. vespertilionis</i>	97
4.2.6	Combined effect of extraction time, temperature and solid to liquid ratio on total phenolic content	100
4.2.7	Modelling of XO inhibitory activity of <i>E. hirta</i> , <i>S. crispa</i> and <i>C. vespertilionis</i>	105
4.2.8	Combined effect of extraction time, temperature and solid to liquid ratio on XO inhibitory activity	107
4.2.9	Optimization and validation of extraction conditions for maximum TFC, TPC and XO inhibitory activity	112

4.3	Cell-based study	113
4.3.1	Effect of plant extracts on the viability of RAW 264.7 macrophage cells	113
4.4	<i>In vivo</i> study of the hyperuricemic rats	117
4.4.1	Acute toxicity study	117
4.4.2	Hypouricemic activity in hyperuricemic rats	121
4.4.3	Liver function test	125
4.4.4	Xanthine oxidase activity in rat liver	128
4.5	Phytochemicals profiling of the plant extracts using liquid chromatography mass spectrometry (LCMS) analysis	130
<b>CHAPTER 5 CONCLUSION AND RECOMMENDATION</b>		<b>135</b>
5.1	Conclusion	135
5.2	Recommendation	137
<b>REFERENCES</b>		<b>138</b>
<b>APPENDICES</b>		<b>178</b>
<b>VITA</b>		<b>205</b>



## LIST OF TABLES

2.1	Sources of purine	13
2.2	The medicinal plants considered to possess anti-gout activity based on <i>in vitro</i> studies	34
2.3	The medicinal plants considered to possess anti-gout activity based on <i>in vivo</i> studies	45
3.1	Experimental design for extraction of plant under different designed conditions	62
3.2	The experimental design of animals (N=72), 6 rats per group	66
4.1	XO inhibitory activity of selected Malaysian medicinal plants.	73
4.2	Pearson's correlation coefficient of TPC and TFC of all the plant extracts with their XO inhibitory activity	80
4.3	Factors and respective low and high values used for screening of significant variables for the extraction of <i>S.crispa</i> , <i>E. hirta</i> and <i>C. vespertilionis</i>	82
4.4	The complete results for TFC, TPC and XO activity of <i>E. hirta</i> , <i>S.crispa</i> and <i>C. vespertilionis</i> based on CCD	87
4.5	Model summary statistics for total flavonoid content of <i>E. hirta</i> , <i>S. crispa</i> and <i>C. vespertilionis</i>	88
4.6	Model summary statistics for total phenolic content of <i>E. hirta</i> , <i>S. crispa</i> and <i>C. vespertilionis</i>	88

4.7	Model summary statistics for XO inhibitory activity of <i>E. hirta</i> , <i>S. crispa</i> and <i>C. vespertilionis</i>	89
4.8	The regression coefficients and results of ANOVA for response surface quadratic model of total flavonoid content of <i>E. hirta</i>	91
4.9	The regression coefficients and results of ANOVA for response surface quadratic model of total flavonoid content of <i>S. crispa</i>	91
4.10	The regression coefficients and results of ANOVA for response surface quadratic model of total flavonoid content of <i>C. vespertilionis</i>	92
4.11	The regression coefficients and results of ANOVA for response surface quadratic model of total phenolic content of <i>E. hirta</i>	98
4.12	The regression coefficients and results of ANOVA for response surface quadratic model of total phenolic content of <i>S. crispa</i>	99
4.13	The regression coefficients and results of ANOVA for response surface quadratic model of total phenolic content of <i>C. vespertilionis</i>	99
4.14	The regression coefficients and results of ANOVA for response surface quadratic model of XO inhibitory activity of <i>E. hirta</i>	106
4.15	The regression coefficients and results of ANOVA for response surface quadratic model of XO inhibitory activity of <i>S. crispa</i>	106
4.16	The regression coefficients and results of ANOVA for response surface quadratic model of XO inhibitory activity of <i>S. crispa</i>	107
4.17	The optimized extraction conditions of <i>E. hirta</i> , <i>S. crispa</i> and <i>C. vespertilionis</i>	112

4.18	Differences between predicted value and experimental value for <i>E. hirta</i> , <i>S. crista</i> and <i>C. vespertilionis</i>	113
4.19	Effects of the aqueous extracts of <i>E. hirta</i> , <i>C. vespertilionis</i> and <i>S. crista</i> on xanthine oxidase activity in the liver of rats	130



## LIST OF FIGURES

2.1	Needle-shaped monosodium crystals seen by light microscopy of synovial fluid in a patient with gout	9
2.2	The chronic gouty tophi of the hands	9
2.3	Four stages of gout	10
2.4	The chemical structure of colchicine	16
2.5	The chemical structure of prednisone	17
2.6	The chemical structure of allopurinol	18
2.7	The chemical structure of febuxostat	19
2.8	The chemical structure of benzbromarone	20
2.9	Pathogenesis of gout	20
2.10	Uric acid structure	22
2.11	Simplified uric acid production and excretion	22
2.12	Uric acid metabolism	23
2.13	The reaction of allopurinol in inhibiting xanthine oxidase activity	24
2.14	<i>Moringa oleifera</i>	25
2.15	<i>Strobilanthes crispus</i>	26
2.16	<i>Artocarpus altilis</i>	27
2.17	<i>Ortosiphon aristatus</i>	28
2.18	<i>Euphorbia hirta</i>	29
2.19	<i>Christia vespertilionis</i>	30
2.20	Central composite design for two and three factors	54
2.21	(a) the cube for Box-behken design (BBD) and three interlocking 22 factorial design (b)	55
3.1	Flowchart of the study design	58

3.2	Acute toxicity chart flow	65
3.3	The flow of the conducted research	67
4.1	Standard calibration curve for gallic acid	76
4.2	Total phenolic content of all the plant extracts	76
4.3	Standard calibration curve for rutin	79
4.4	Total flavonoid content of all the plant extracts	79
4.5	Pareto chart for the screening of parameters for plant extraction of <i>E. hirta</i> towards the responses	83
4.6	Pareto chart for the screening of parameters for plant extraction of <i>S. crispa</i> towards the responses	84
4.7	Pareto chart for the screening of parameters for plant extraction of <i>C. vespertilionis</i> towards the responses	85
4.8	Response surface plots of <i>E. hirta</i> on total flavonoid content	94
4.9	Response surface plots of <i>C. vespertilionis</i> on total flavonoid content	95
4.10	Response surface plots of <i>S. crispa</i> on total flavonoid content	96
4.11	Response surface plots of <i>E. hirta</i> on total phenolic content	102
4.12	Response surface plots of <i>S. crispa</i> on total phenolic content	103
4.13	Response surface plots of <i>C. vespertilionis</i> on total phenolic content	104
4.14	Response surface plots of <i>E. hirta</i> leaves on XO inhibitory activity	109
4.15	Response surface plots of <i>S. crispa</i> leaves on XO inhibitory activity	110

4.16	Response surface plots of <i>C. vespertilionis</i> leaves on XO inhibitory activity	111
4.17	Effects of treatment with <i>E. hirta</i> at various concentrations on cell viability of RAW 264.7 cells	116
4.18	Effects of treatment with <i>S. crispa</i> at various concentrations on cell viability of RAW 264.7 cells	116
4.19	Effects of treatment with <i>C. vespertilionis</i> at various concentrations on cell viability of RAW 264.7 cells	117
4.20	Mean body weight of male rats receiving <i>E. hirta</i> , <i>C. vespertilionis</i> and <i>S. crispa</i> extracts for 14 days (first set)	119
4.21	Mean body weight of male rats receiving <i>E. hirta</i> , <i>C. vespertilionis</i> and <i>S. crispa</i> extracts for 14 days (second set)	120
4.22	Mean of weights of organs of male rats receiving <i>E. hirta</i> , <i>C. vespertilionis</i> and <i>S. crispa</i> extracts for 14 days (first set)	120
4.23	Mean of weights of organs of male rats receiving <i>E. hirta</i> , <i>C. vespertilionis</i> and <i>S. crispa</i> extracts for 14 days (second set)	121
4.24	Effects of the aqueous extracts of <i>E. hirta</i> , <i>C. vespertilionis</i> and <i>S. crispa</i> on serum urate levels in rats treated with the uricase inhibitor potassium oxonate	124
4.25	AST levels in rats administered with <i>C. vespertilionis</i> , <i>E. hirta</i> and <i>S. crispa</i> extracts	127
4.26	ALT levels in rats administered with <i>C. vespertilionis</i> , <i>E. hirta</i> and <i>S. crispa</i> extracts	127
4.27	GGT levels in rats administered with <i>C. vespertilionis</i> , <i>E. hirta</i> and <i>S. crispa</i> extracts	128

4.28	LCMS chromatogram, acquired in the negative-ion mode, from an extract of the <i>E. hirta</i>	133
4.29	LCMS chromatogram, acquired in the negative-ion mode, from an extract of the <i>S. crispa</i>	134
4.30	LCMS chromatogram, acquired in the negative-ion mode, from an extract of the <i>C. vespertilionis</i>	134



## LIST OF SYMBOLS AND ABBREVIATIONS

°C	-	Degree celcius
g	-	Gram
h	-	Hour
mg	-	Milligram
min	-	Minute
ml	-	Milliliter
µg	-	Microgram
mg/dl	-	Milligram per deciliter
mg/g	-	Milligram per g
µg/ml	-	Microgram per milliliter
µmol/l	-	Micromol per liter
µl	-	Microliter
<i>r</i>	-	Pearson correlation coefficient
<i>r</i> <sup>2</sup>	-	Coefficient of determination
2FI	-	Two factor interaction
ACR	-	American College of Rheumatologist
AlCl <sub>3</sub> .6H <sub>2</sub> O	-	Aluminium chloride hexahydrate
ALT	-	Alanine aminotransferase
ANOVA	-	Analysis of variance
AST	-	Aspartate Aminotransferase
ATP	-	Adenosine triphosphate
BBD	-	Box-Behnken design
BSR	-	British Society of Rheumatologist
CCD	-	Central composite design
CID	-	Collision dissociation
CMC	-	Carboxymethyl cellulose
CO <sub>2</sub>	-	Carbon dioxide



COXIBS	-	COX-2 inhibitors
COX-2	-	Cyclooxygenase 2
DMSO	-	Dimethylsulfoxide
DNA	-	Deoxyribunucleic acid
EDTA	-	Ethylenediaminetetraacetic acid
EPP	-	Entry Point Project
ETP	-	Economic transformation program
EULAR	-	European League Against Rheumatism
FAD	-	Flavin adenine dinucleotide
FDA	-	Food and Drug Administration
GAE	-	Gallic acid equivalents
GGT	-	Gamma-glutamyl transferase
GLUT9	-	Glucose transporter 9
GNI	-	Gross national income
HCl	-	Hydrochloric acid
HPLC	-	High performance liquid chromatography
IL-1 $\beta$	-	Interleukin-1 beta
IL-6	-	Interleukin-6
iNOS	-	Inducible nitric oxide synthase
ISO	-	International Organization for Standardization
KCl	-	Potassium chloride
LCMS	-	Liquid chromatography mass spectrometry
MCP-1	-	Monocyte chemoattractant protein-1
MPC-SNEDDS	-	Self-nanoemulsifying drug delivery system based on morin-phospholipid complex
MRM	-	Multiple reaction monitoring
MSU	-	Monosodium urate
MTT	-	3-(4, 5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide
NA	-	Not available
NaNO <sub>2</sub>	-	Sodium nitrite
NaOH	-	Sodium hydroxide
NIAMS	-	National Institute of Arthritis and Musculoskeletal and Skin Diseases

NKEA	-	National Key Economic Area
NLRP3	-	Nod-like receptor family pyrin domain containing 3
NMR	-	Nuclear magnetic resonance
NO	-	Nitric oxide
NSAID	-	Non-steroidal anti-inflammatory drug
OCT	-	Organic cation transporter
OCTN	-	Carnitine transporter
OECD	-	Organization for Economic Cooperation and Development
OH	-	Hydroxyl group
PBS	-	Phosphate buffered saline
PEMANDU	-	Performance Management & Delivery unit
PGE2	-	Prostaglandin E <sub>2</sub>
PO	-	Potassium oxonate
QE	-	Quercetin equivalents
RE	-	Rutin equivalents
RNA	-	Ribonucleic acid
RPMI	-	Roswell Park Application of Technology
RSM	-	Response surface methodology
SEM	-	Standard error of mean
SLC2A9	-	Solute carrier family 2 member 9
SPSS	-	Statistical Package for the Social Sciences
TFC	-	Total flavonoid content
TNF- $\alpha$	-	Tumor necrosis factor-alpha
TNF- $\beta$	-	Tumor necrosis factor-beta
TPC	-	Total phenolic content
URAT1	-	Urate transporter 1
UV	-	Ultraviolet
UV-VIS	-	Ultraviolet-visible
VEGF	-	Vascular endothelial growth factor
WHO	-	World Health Organization
XO	-	Xanthine oxidase

## LIST OF APPENDICES

APPENDIX	TITLE	PAGE
A	Letter of confirmation of plant species	178
B	Cytotoxicity study of RAW264.7 cells	179
C	Approval letter for ethics	180
D	<i>In vivo</i> study on hyperuricemic rats	181
E	ANOVA test for the XO inhibitory activity of the plants	182
F	Total phenolic and flavonoid contents of the six plant extracts	183
G	Pearson's correlation analysis between the phytochemical contents and anti-gout activity	184
H	Normal plot distribution of <i>E. hirta</i> , <i>S. crispa</i> and <i>C. vespertilionis</i>	185
I	Cytotoxic effect of <i>E. hirta</i> , <i>S. crispa</i> and <i>C. vespertilionis</i> on mouse macrophage RAW264.7 cell	188
J	Percentage of cell viability (%) treated with <i>E. hirta</i> , <i>C. vespertilionis</i> and <i>S. crispa</i> aqueous extracts for 24 h	191
K	Body weight (g) of rats in acute toxicity test	192
L	Organ weight (g) of rats receiving single dose of <i>S. crispa</i> , <i>E. hirta</i> and <i>C. vespertilionis</i>	193
M	Effects of the aqueous extracts of <i>E. hirta</i> , <i>C. vespertilionis</i> and <i>S. crispa</i> on serum urate	

	levels in rats treated with the uricase inhibitor, potassium oxonate.	194
N	Effects of the aqueous extracts of <i>E. hirta</i> , <i>C. vespertilionis</i> and <i>S. crispa</i> on AST, ALT and GGT levels in the liver of rats	195
O	Chemical constituents present in the <i>E. hirta</i> <i>S. crispa</i> and <i>C. vespertilionis</i> aqueous extract	196
P	Mass spectrum of <i>E. hirta</i> , <i>S. crispa</i> and <i>C. vespertilionis</i> extract in negative ion mode	197



## REFERENCES

- Abdellatif, A. A., & Elkhaili, N. (2014). Management of gouty arthritis in patients with chronic kidney disease. *American Journal of Therapeutics*, 21(6), 523-534.
- Abd Aziz, S. M., Low, C. N., Chai, L. C., Abd Razak, S. S. N., Selamat, J., Son, R., Sarker, M. Z. I. & Khatib, A. (2011). Screening of selected Malaysian plants against several food borne pathogen bacteria. *International Food Research Journal*, 18(3), 1195- 1201.
- Abhishek, A., Valdes, A. M., Jenkins, W., Zhang, W., & Doherty, M. (2017). Triggers of acute attacks of gout, does age of gout onset matter? A primary care based cross-sectional study. *PloS one*, 12(10), e0186096.
- Abu-Gharbieh, E., Shehab, N. G., Almasri, I. M., & Bustanji, Y. (2018). Antihyperuricemic and xanthine oxidase inhibitory activities of *Tribulus arabicus* and its isolated compound, ursolic acid: *in vitro* and *in vivo* investigation and docking simulations. *PloS one*, 13(8), e0202572.
- Acero, N., Gradillas, A., Beltran, M., García, A., & Mingarro, D. M. (2019). Comparison of phenolic compounds profile and antioxidant properties of different sweet cherry (*Prunus avium* L.) varieties. *Food Chemistry*, 279, 260-271.
- Afrizal, I.S.P. (2008). *Analytical, bioactivity and stability studies on Strobilanthes crispus L. Bremek and Sonchus Arvensis L. extracts*. Universiti Sains Malaysia: Thesis Ph.D.
- Agata, I., Hatano, T., Nakaya, Y., Sugaya, T., Nishibe, S., Yoshida, T., & Okuda, T. 1(1991). Tannins and related polyphenols of Euphorbiaceous Plants. VIII. Eumaculin A and Eusupinin A, and accompanying polyphenols from *Euphorbia maculata* L. *Chemical and Pharmaceutical Bulletin*, 39(4), 881-883.

- Akowuah, G. A., Zhari, I., Norhayati, I., Sadikun, A., & Khamsah, S. M. (2004). Sinensetin, eupatorin, 30 -hydroxy-5, 6, 7, 40 -tetramethoxyflavone and rosmarinic acid contents and antioxidative effect of *Orthosiphon stamineus* from Malaysia. *Food Chemistry*, 87(4), 559-566.
- Akowuah, G. A., Ismail, Z., Norhayati, I., & Sadikun, A. (2005). The effects of different extraction solvents of varying polarities on polyphenols of *Orthosiphon stamineus* and evaluation of the free radical-scavenging activity. *Food Chemistry*, 93(2), 311-317.
- Al-Azzawie, H. F., & Abd S. A. (2015). Effect of some plant extracts on serum uric acid levels and xanthine oxidase activity *in vitro* and in oxonate-induced hyperuricemic rats. *European Journal of Pharmaceutical and Medical Research*, 2(6), 55-61.
- Al-Farsi, M. A., & Lee, C. Y. (2008). Optimization of phenolics and dietary fibre extraction from date seeds. *Food Chemistry*, 108(3), 977-985.
- Almatar, M., & Rahmat, Z. (2014). Identifying the developmental stages and optimizing the sample preparation for anatomical study of *Orthosiphon stamineus*. *Journal of Applied Pharmaceutical Science*, 4(3), 66-74.
- Alsultane, I. R., Ewadh, M. J., & Mohammed, M. F. (2014). Novel natural anti gout medication extract from *Momordica charantia*. *Journal of Natural Sciences Research*, 4(17), 16-23.
- Alvarez-Lario, B., & Macarron-Vicente, J. (2011). Is there anything good in uric acid?. *QJM: An International Journal of Medicine*, 104(12), 1015-1024.
- Amid, M., Manap, A., Yazid, M., & Zohdi, N. (2014). Optimization of processing parameters for extraction of amylase enzyme from Dragon (*Hylocereus polyrhizus*) peel using response surface methodology. *The Scientific World Journal*, 2014, 1-12.
- Amna, O. F., Nooraain, H., Noriham, A., Azizah, A. H., & Husna, R. N. (2013). Acute and oral subacute toxicity study of ethanolic extract of *Cosmos caudatus* leaf in Sprague Dawley rats. *International Journal of Bioscience, Biochemistry and Bioinformatics*, 3(4), 301- 305.
- Anbari, K. K. (2014). *Gout causes and risk factors*. Retrieved February 25, 2019, from <https://www.arthritis-health.com/types/gout/gout-causes-and-risk-factors>

- Antón, F. M., Puig, J. G., Ramos, T., Gonzalez, P., & Ordas, J. (1986). Sex differences in uric acid metabolism in adults: evidence for a lack of influence of estradiol-17 $\beta$  (E2) on the renal handling of urate. *Metabolism*, 35(4), 343-348.
- Anwar, F., Latif, S., Ashraf, M., & Gilani, A. H. (2007). *Moringa oleifera*: a food plant with multiple medicinal uses. *Phytotherapy*, 21(1), 17-25.
- Apaya, K. L., & Chichioco-Hern, C. L. (2011). Xanthine oxidase inhibition of selected Philippine medicinal plants. *Journal of Medicinal Plants Research*, 5(2), 289-292.
- Arafat, O. M., Tham, S. Y., Sadikun, A., Zhari, I., Haughton, P. J., & Asmawi, M. Z., (2008). Studies on diuretic and hypouricemic effects of *Orthosiphon stamineus* methanol extracts in rats. *Journal of Ethnopharmacology*, 118(3), 354-360.
- Arromdee, E., Michet, C.J., Crowson, C.S., O'Fallon, W.M., & Gabriel, S.E. (2002). Epidemiology of gout: is the incidence rising? *The Journal of Rheumatology*, 29(11), 2403-2406.
- Auletta, C. S. (1995). *Acute, subchronic and chronic toxicology*. London: CRC Press.
- Awale, S., Tezuka, Y., Banskota, A. H., & Kadota, S. (2002). Siphonols A-E: Novel nitric oxide inhibitors from *Orthosiphon stamineus* of Indonesia. *Bioorganic and Medicinal Chemistry Letters*, 13(1), 31-35.
- Azahar, N. F., Gani, S. S. A., & Mokhtar, N. F. M. (2017). Optimization of phenolics and flavonoids extraction conditions of *Curcuma zedoaria* leaves using response surface methodology. *Chemistry Central Journal*, 11(1), 96.
- Azmi, S. M. N., Jamal, P., & Amid, A. (2012). Xanthine oxidase inhibitory activity from potential Malaysian medicinal plant as remedies for gout. *International Food Research Journal*, 19(1), 59-66.
- Bailey, P. S., & Bailey, C. A. (2000). *Organic chemistry: a brief survey of concepts and applications*. Upper Saddle River: Prentice Hall.
- Basma, A. A., Zakaria, Z., Latha, L. Y., & Sasidharan, S. (2011). Antioxidant activity and phytochemical screening of the methanol extracts of *Euphorbia hirta* L. *Asian Pacific Journal of Tropical Medicine*, 4(5), 386-390.
- Becker, M. A., Schumacher Jr, H. R., Wortmann, R. L., MacDonald, P. A., Eustace, D., Palo, W. A., Streit, J., & Joseph-Ridge, N. (2005). Febuxostat compared with allopurinol in patients with hyperuricemia and gout. *New England Journal of Medicine*, 353(23), 2450-2461.



- Becker, M. A., Schumacher, H. R., Espinoza, L. R., Wells, A. F., MacDonald, P., Lloyd, E., & Lademacher, C. (2010). The urate-lowering efficacy and safety of febuxostat in the treatment of the hyperuricemia of gout: the CONFIRMS trial. *Arthritis Research & Therapy*, 12(2), R63.
- Bello, I., Bakkouri, A. S., Tabana, Y. M., Al-Hindi, B., Al-Mansoub, M. A., Mahmud, R., & Asmawi, M. Z. (2016). Acute and sub-acute toxicity evaluation of the methanolic extract of *Alstonia scholaris* stem bark. *Medical Sciences*, 4(1), 4.
- Belwal, T., Dhyani, P., Bhatt, I. D., Rawal, R. S., & Pande, V. (2016). Optimization extraction conditions for improving phenolic content and antioxidant activity in *Berberis asiatica* fruits using response surface methodology (RSM). *Food Chemistry*, 207, 115-124.
- Ben Mohamed Maoulainine, L., Jelassi, A., Hassen, I., & Ould Mohamed Salem Ould Boukhari, A. (2012). Antioxidant proprieties of methanolic and ethanolic extracts of *Euphorbia helioscopia* (L.) aerial parts. *International Food Research Journal*, 19(3), 1125-1130.
- Bezerra, M. A., Santelli, R. E., Oliveira, E. P., Villar, L. S., & Escalera, L. A. (2008). Response surface methodology (RSM) as a tool for optimization in analytical chemistry. *Talanta*, 76(5), 965-977.
- Bimakr, M., Rahman, R. A., Taip, F. S., Ganjloo, A., Salleh, L. M., Selamat, J., Hamid, A., & Zaidul, I. S. M. (2011). Comparison of different extraction methods for the extraction of major bioactive flavonoid compounds from spearmint (*Mentha spicata* L.) leaves. *Food and Bioproducts Processing*, 89(1), 67-72.
- Borges, F., Fernandes, E., & Roleira, F. (2002). Progress towards the discovery of xanthine oxidase inhibitors. *Current Medicinal Chemistry*, 9(2), 195-217.
- Box, G. E., & Behnken, D. W. (1960). Some new three level designs for the study of quantitative variables. *Technometrics*, 2(4), 455-475.
- Box, G. E., Hunter, J. S., & Hunter, W. G. (2005). *Statistics for experimenters: design, innovation, and discovery*. New York: Wiley-Interscience.
- Brach, A. R., & Song, H. (2006). eFloras: New directions for online floras exemplified by the Flora of China Project. *Taxon*, 55(1), 188-192.
- Bruns, R. E., Scarminio, I. S., & de Barros Neto, B. (2006). *Statistical design-chemometrics*. Amsterdam: Elsevier.
- Buckley, H. R. (2011). Epidemiology of gout: Perspectives from the past. *Current Rheumatology Reviews*, 7(2), 106-113.



- Buenz, E.J., Schneppele, D.J., Bauer, B.A., Elkin, P.L., Riddle, J.M., & Motley, T.J. (2004). Techniques: Bioprospecting historical herbal texts by hunting for new leads in old tomes. *Trends in Pharmacological Sciences*, 25(9), 494-498.
- Bunawan, H., Bunawan, S. N., & Baharum, S. N. (2015). The red butterfly wing (*Christia vespertilionis*): a promising cancer cure in Malaysia. *International Journal of Pharmacy and Pharmaceutical Sciences*, 7(8), 1.
- Burk, D. R., Senechal-Willis, P., Lopez, L. C., Hogue, B. G., & Daskalova, S. M. (2009). Suppression of lipopolysaccharide-induced inflammatory responses in RAW 264.7 murine macrophages by aqueous extract of *Clinopodium vulgare* L. (Lamiaceae). *Journal of Ethnopharmacology*, 126(3), 397-405.
- Burns, C. M., & Wortmann, R. L. (2012). Latest evidence on gout management: what the clinician needs to know. *Therapeutic Advances in Chronic Disease*, 3(6), 271-286.
- Bustanji, Y., Hudaib, M., Tawaha, K., Mohammad, M. K., Almasri, I., Hamed, S., & Oran, S. (2011). *In vitro* xanthine oxidase inhibition by selected Jordanian medicinal plants. *Jordan Journal of Pharmaceutical Sciences*, 4(1), 49-56.
- Bystrická, J., Vollmannová, A., & Margitanová, E. (2010). Dynamics of polyphenolics formation in different plant parts and different growth phases of selected buckwheat cultivars. *Acta Agriculturae Slovenica*, 95(3), 225- 229.
- Cacace, J. E., & Mazza, G. (2003). Mass transfer process during extraction of phenolic compounds from milled berries. *Journal of Food Engineering*, 59(4), 379-389.
- Cacciola, F., Delmonte, P., Jaworska, K., Dugo, P., Mondello, L., & Rader, J. I. (2011). Employing ultra high pressure liquid chromatography as the second dimension in a comprehensive two-dimensional system for analysis of *Stevia rebaudiana* extracts. *Journal of Chromatography A*, 1218(15), 2012-2018.
- Cai, Z., Lee, F. S. C., Wang, X. R., & Yu, W. J. (2002). A capsule review of recent studies on the application of mass spectrometry in the analysis of Chinese medicinal herbs. *Journal of Mass Spectrometry*, 37(10), 1013-1024.
- Cai, Y., Luo, Q., Sun, M., & Corke, H. (2004). Antioxidant activity and phenolic compounds of 112 Chinese medicinal plants associated with anticancer. *Life Science*, 74(17), 2157-2184.
- Carley, K. M., Kamneva, N. Y., & Reminga, J. (2004). Response surface methodology (No. CMU-ISRI-04-136). USA: Carnegie-Mellon University.

- Carro, M. D., Falkenstein, E., Blemings, K. P., & Klandorf, H. (2009). Determination of xanthine oxidoreductase activity in broilers: Effect of pH and temperature of the assay and distribution in tissues. *Poultry Science*, 88(11), 2406-2414.
- Caulfield, M. J., Munroe, P. B., O'Neill, D., Witkowska, K., Charchar, F. J., Doblado, M., Evans, S., Eyheramendy, S., Onipinla, A., Howard, P., & Shaw-Hawkins, S. (2008). SLC2A9 is a high-capacity urate transporter in humans. *PLoS Medicine*, 5(10), e197.
- Chan, W. S., Wen, P. C., & Chiang, H. C. (1995). Structure-activity relationship of caffeic acid analogues on xanthine oxidase inhibition. *Anticancer Research*, 15(3), 703-707.
- Chan, E. W. C., Lim, Y. Y., Chong, K. L., Tan, J. B. L., & Wong, S. K. (2010). Antioxidant properties of tropical and temperate herbal teas. *Journal of Food Composition and Analysis*, 23(2), 185-189.
- Chang, H. Y., Pan, W. H., Yeh, W. T., & Tsai, K. S. (2001). Hyperuricemia and gout in Taiwan: results from the Nutritional and Health Survey in Taiwan (1993-96). *The Journal of Rheumatology*, 28(7), 1640-1646.
- Chang, Y. C., Lee, F. W., Chen, C. S., Huang, S. T., Tsai, S. H., Huang, S. H., & Lin, C. M. (2007). Structure-activity relationship of C6-C3 phenylpropanoids on xanthine oxidase-inhibiting and free radical-scavenging activities. *Free Radical Biology and Medicine*, 43(11), 1541-1551.
- Chao, J., & Terkeltaub, R. (2009). A critical reappraisal of allopurinol dosing, safety, and efficacy for hyperuricemia in gout. *Current Rheumatology Reports*, 11(2), 135-140.
- Chen, L., Yin, H., Lan, Z., Ma, S., Zhang, C., Yang, Z., Li, P., & Lin, B. (2011). Anti-hyperuricemic and nephroprotective effects of *Smilax china* L. *Journal of Ethnopharmacology*, 135(2), 399-405.
- Chen, J. H., Yeh, W. T., Chuang, S. Y., Wu, Y. Y., & Pan, W. H. (2012). Gender-specific risk factors for incident gout: a prospective cohort study. *Clinical Rheumatology*, 31(2), 239-245.
- Cheng, X., & Changgui, L. (2015). The principles of gout therapy. *Gout and Hyperuricemia*, 2(1), 15-23.

- Cheng, L. C., Murugaiyah, V., & Chan, K. L. (2015). Flavonoids and phenylethanoid glycosides from *Lippia nodiflora* as promising antihyperuricemic agents and elucidation of their mechanism of action. *Journal of Ethnopharmacology*, 176, 485-493.
- Choi, H. K., Atkinson, K., Karlson, E. W., Willett, W., & Curhan, G. (2004). Purine-rich foods, dairy and protein intake, and the risk of gout in men. *New England Journal of Medicine*, 350(11), 1093-1103.
- Choi, H. K., Mount, D. B., & Reginato, A. M. (2005a). Pathogenesis of gout. *Annals of Internal Medicine*, 143(7), 499-516.
- Choi, H. K., & Curhan, G. (2005b). Gout: epidemiology and lifestyle choices. *Current Opinion in Rheumatology*, 17(3), 341-345.
- Choi, H. K., Atkinson, K., Karlson, E. W., & Curhan, G. (2005c). Obesity, weight change, hypertension, diuretic use, and risk of gout in men: the health professionals follow-up study. *Archives of Internal Medicine*, 165(7), 742-748.
- Choi, H. K., & Ford, E. S. (2007). Prevalence of the metabolic syndrome in individuals with hyperuricemia. *The American Journal of Medicine*, 120(5), 442-447.
- Chuang, S. Y., Lee, S. C., Hsieh, Y. T., & Pan, W. H. (2011). Trends in hyperuricemia and gout prevalence: Nutrition and Health Survey in Taiwan from 1993-1996 to 2005-2008. *Asia Pacific Journal of Clinical Nutrition*, 20(2), 301-308.
- Conway, N., & Schwartz, S. (2009). Diagnosis and management of acute gout. *Rhode Island Medical Journal*, 92(11), 356-358.
- Coppin, J. P., Xu, Y., Chen, H., Pan, M. H., Ho, C. T., Juliani, R., Simon, J. E., & Wu, Q. (2013). Determination of flavonoids by LC/MS and anti-inflammatory activity in *Moringa oleifera*. *Journal of Functional Foods*, 5(4), 1892-1899.
- Cos, P., Ying, L., Calomme, M., Hu, J. P., Cimanga, K., Van Poel, B., Pieters, L., Vlietinck, A. J., & Berghe, D. V. (1998). Structure- activity relationship and classification of flavonoids as inhibitors of xanthine oxidase and superoxide scavengers. *Journal of Natural Products*, 61(1), 71-76.
- Costa, A. R., Freitas, L. A., Mendiola, J., & Ibáñez, E. (2015). *Copaifera langsdorffii* supercritical fluid extraction: Chemical and functional characterization by LC/MS and *in vitro* assays. *The Journal of Supercritical Fluids*, 100, 86-96.
- Cornell, J. A. (2011). *Experiments with mixtures: designs, models, and the analysis of mixture data*. New York: Wiley.

- Cucuianu, M., & Brudasca, I. (2012). Gout, hyperuricemia and the metabolic syndrome. *Romanian Journal of Laboratory Medicine*, 20(3), 199-206.
- Dai, J., & Mumper, R. J. (2010). Plant phenolics: extraction, analysis and their antioxidant and anticancer properties. *Molecules*, 15(10), 7313-7352.
- Dandekar, R., Fegade, B., & Naik, A. (2015). Evaluation of anti-inflammatory activity of alcohol and aqueous extract of *Epiphyllum oxypetalum* leaves. *World Journal of Pharmacy and Pharmaceutical Sciences*, 4(7), 851-858.
- Davis, J. M., Murphy, E. A., Carmichael, M. D., & Davis, B. (2009). Quercetin increases brain and muscle mitochondrial biogenesis and exercise tolerance. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, 296(4), R1071-R1077.
- Deng, B., Liu, Z., & Zou, Z. (2019). Optimization of microwave-assisted extraction saponins from *Sapindus mukorossi* pericarps and an evaluation of their inhibitory activity on xanthine oxidase. *Journal of Chemistry*, 2019, 1-19.
- Derle, D. V., Gujar, K. N., & Sagar, B. S. H. (2006). Adverse effects associated with the use of nonsteroidal antiinflammatory drugs: An overview. *Indian Journal of Pharmaceutical Sciences*, 68(4), 409- 414.
- Dessein, P. H., Shipton, E. A., Stanwix, A. E., Joffe, B. I., & Ramokgadi, J. (2000). Beneficial effects of weight loss associated with moderate calorie/carbohydrate restriction, and increased proportional intake of protein and unsaturated fat on serum urate and lipoprotein levels in gout: a pilot study. *Annals of the Rheumatic Diseases*, 59(7), 539-543.
- De Souza, A. W., Fernandes, V., & Ferrari, A. J. (2005). Female gout: clinical and laboratory features. *The Journal of Rheumatology*, 32(11), 2186-2188.
- Dhanasekar, C., Kalaiselvan, S., & Rasool, M. (2015). Morin, a bioflavonoid suppresses monosodium urate crystal-induced inflammatory immune response in RAW 264.7 macrophages through the inhibition of inflammatory mediators, intracellular ROS levels and NF- $\kappa$ B activation. *PLoS One*, 10(12), e0145093.
- Di Giovine, F. S., Malawista, S. E., Nuki, G., & Duff, G. W. (1987). Interleukin 1 (IL 1) as a mediator of crystal arthritis. Stimulation of T cell and synovial fibroblast mitogenesis by urate crystal-induced IL 1. *The Journal of Immunology*, 138(10), 3213-3218.

- Do, Q. D., Angkawijaya, A. E., Tran-Nguyen, P. L., Huynh, L. H., Soetaredjo, F. E., Ismadji, S., & Ju, Y. H. (2014). Effect of extraction solvent on total phenol content, total flavonoid content, and antioxidant activity of *Limnophila aromatica*. *Journal of Food and Drug Analysis*, 22(3), 296-302.
- Dybing, E., Doe, J., Groten, J., Kleiner, J., O'brien, J., Renwick, A. G., Schlatter, J., Steinberg, P., Tritscher, A., Walker, R., & Younes, M. (2002). Hazard characterisation of chemicals in food and diet: dose response, mechanisms and extrapolation issues. *Food and Chemical Toxicology*, 40(2-3), 237-282.
- Eashwary, M., & Hussein, H. (2006). Disease of Kings-clinical characteristics at two tertiary referral centers in Malaysia. *APLAR Journal of Rheumatology*, 9, A21-A278.
- Edirs, S., Turak, A., Numonov, S., Xin, X., & Aisa, H. A. (2017). Optimization of extraction process for antidiabetic and antioxidant activities of Kursi Wufarikun Ziyabit using response surface methodology and quantitative analysis of main components. *Evidence-Based Complementary and Alternative Medicine*, 2017, 1-14.
- Eggebeen, A.T. (2007). Gout: an update. *American Family Physician*, 76(6), 801-808.
- El-Boghdady, N. A. (2011). Protective effect of ellagic acid and pumpkin seed oil against methotrexate-induced small intestine damage in rats. *Indian Journal of Biochemistry & Biophysics*, 48(6), 380-387.
- El-Rahman, H.S., & Abd-Elhak, N. A. (2015). Xanthine oxidase inhibitory activity and antigout of celery leek parsley and molokhia. *Advances in Biochemistry*, 3(4), 40-50.
- Elijah, J. P., Young, U. C., Chuma, E., Chinelo, N. C., Onyiaoziri, O. J., Ogara, A., Obiageli, N. F., Vincent, O., Oteri, E. J., & Emmanuela, U. N. (2017). Biochemical effects of the aqueous extract of *Hibiscus sabdariffa* on liver marker enzymes and lipid profiles in acetaminophen-challenged rats. *Research Journal of Medicinal Plants*, 11(2), 48-54.
- Enomoto, A., Kimura, H., Chairoungdua, A., Shigeta, Y., Jutabha, P., Cha, S. H., Hosoyamada, M., Takeda, M., Sekine, T., Igarashi, T., & Matsuo, H. (2002). Molecular identification of a renal urate-anion exchanger that regulates blood urate levels. *Nature*, 417(6887), 447-452.



- Eugster, P. J., Guillarme, D., Rudaz, S., Veuthey, J. L., Carrupt, P. A., & Wolfender, J. L. (2011). Ultra high pressure liquid chromatography for crude plant extract profiling. *Journal of AOAC International*, 94(1), 51-70.
- Fadzureena, J., Mazura, M. P., Adiana, M. A., & Hani, I. B. (2013). An investigation into the inhibitory effect of *Senna alata* L. leaf extract as well as its isolated compound on xanthine oxidase assay. *Proceedings of the Conference on Forestry and Forest Products Research*. Kuala Lumpur, Malaysia. 262.
- Fahey, J.W. (2005). *Moringa oleifera*: A review of the medical evidence for its nutritional, therapeutic, and prophylactic properties part 1. *Trees for Life Journal*, 1, 5-15.
- Falasca, G. F. (2006). Metabolic diseases: gout. *Clinics in Dermatology*, 24(6), 498-508.
- Falodun, A., Ali, S., Quadir, I. M., & Choudhary, I. M. (2008). Phytochemical and biological investigation of chloroform and ethylacetate fractions of *Euphorbia heterophylla* leaf (Euphorbiaceae). *Journal of Medicinal Plants Research*, 2(12), 365-369.
- Fariza, I., Fadzureena, J., Zunoliza, A., Chuah, A. L., Pin, K. Y., & Adawiah, I. (2012). Anti-inflammatory activity of the major compound from methanol extract of *Phaleria macrocarpa* leaves. *Journal of Applied Sciences*, 12(11), 1195-1198.
- Fathallah, N., Salem, C. B., Slim, R., Kaabia, N., Letaief, A., & Bouraoui, K. (2010). Fatal allopurinol-induced hypersensitivity syndrome associated with pancreatic abnormalities. *JCR: Journal of Clinical Rheumatology*, 16(4), 170-171.
- Fejes, S.Z., Blazovics, A., Lemberkovics, E., Petri, G., Szöke, E., & Kery, A. (2000). Free radical scavenging and membrane protective effects of methanol extracts from *Anthriscus cerefolium* L.(Hoffm.) and *Petroselinum crispum* (Mill.) Nym. ex AW Hill. *Phytotherapy Research*, 14(5), 362-365.
- Ferreira, S. L. C., Bruns, R. E., da Silva, E. G. P., dos Santos, W. N. L., Quintella, C. M., David, J. M., de Andrade, J. B., Breitzkreitz, M. C., Jardim, I. C. F. S., & Neto, B. B. (2007). Statistical designs and response surface techniques for the optimization of chromatographic systems. *Journal of Chromatography A*, 1158(1-2), 2-14.

- Flemmig, J., Kuchta, K., Arnhold, J., & Rauwald, H. W. (2011). *Olea europaea* leaf (Ph. Eur.) extract as well as several of its isolated phenolics inhibit the gout-related enzyme xanthine oxidase. *Phytomedicine*, 18(7), 561-566.
- Galvez, J., Zarzuelo, A., Crespo, M. E., Lorente, M. D., Ocete, M. A., & Jimenez, J. (1993). Antidiarrhoeic activity of *Euphorbia hirta* extract and isolation of an active flavonoid constituent. *Planta Medica*, 59(04), 333-336.
- Ge, Q., Huang, J., Mao, J. W., Gong, J. Y., Zhou, Y. F., & Huang, J. X. (2014). Optimization of total polysaccharide extraction from *Herba Lophatheri* using RSM and antioxidant activities. *International Journal of Biological Macromolecules*, 67, 37-42.
- Ghasemzadeh, A., Jaafar, H. Z., & Rahmat, A. (2015). Phytochemical constituents and biological activities of different extracts of *Strobilanthes crispus* (L.) Bremek leaves grown in different locations of Malaysia. *BMC Complementary and Alternative Medicine*, 15(1), 422.
- Ghitescu, R. E., Volf, I., Carausu, C., Bühlmann, A. M., Gilca, I. A., & Popa, V. I. (2015). Optimization of ultrasound-assisted extraction of polyphenols from spruce wood bark. *Ultrasonics Sonochemistry*, 22, 535-541.
- Gibson, T., Waterworth, R., Hatfield, P., Robinson, G., & Bremner, K. (1984). Hyperuricaemia, gout and kidney function in New Zealand Maori men. *Rheumatology*, 23(4), 276-282.
- Gide, M., Sharma, P., Saudagar, R., & Shrivastava, B. (2014). Method development and validation for determination of febuxostat from spiked human plasma using RP-HPLC with UV detection. *Chromatography Research International*, 2014, 1-5.
- Gińdzieńska-Sieśkiewicz, E., Sierakowski, S., Domysławska, I., & Sulik, A. (2010). Dna moczanowa—aktualne spojrzenie na diagnostykę i leczenie. *Rheumatology*, 48(6), 425-428.
- Goh, K. L. (2004). *Malaysian herbs*. Klang: Goh Kong Ling.
- Golden, K. D., & Williams, O. J. (2001). Amino acid, fatty acid, and carbohydrate content of *Artocarpus altilis* (breadfruit). *Journal of Chromatographic Science*, 39(6), 243-250.
- Gopalakrishnanb, L., Doriya, K., & Kumara, D.S. (2016). *Moringa oleifera*: A review on nutritive importance and its medicinal application. *Food Science and Human Wellness*, 5(2), 49-56.

- Gopi, K., Anbarasu, K., Renu, K., Jayanthi, S., Vishwanath, B. S., & Jayaraman, G. (2016). Quercetin-3-O-rhamnoside from *Euphorbia hirta* protects against snake venom induced toxicity. *Biochimica et Biophysica Acta (BBA)-General Subjects*, 1860(7), 1528-1540.
- Gothai, S., Vijayarathna, S., Chen, Y., Lai, N. S., Wahab, H. A., Firdaus, H., Sreeramanan, S., & Sasidharana, S. (2019). *In vitro* and *in vivo*-scientific evaluation on cytotoxicity and genotoxicity of traditional medicinal plant *Couroupita guianensis* Aubl. flower. *Pharmacologyonline*, 2, 24-38.
- Gottlieb, O. R., & Borin, M. R. D. (2000). Medicinal products: regulation of biosynthesis in space and time. *Memórias do Instituto Oswaldo Cruz*, 95(1), 115-120.
- Grata, E., Boccard, J., Guillarme, D., Glauser, G., Carrupt, P. A., Farmer, E. E., Wolfender, J. L., & Rudaz, S. (2008). UPLC–TOF-MS for plant metabolomics: a sequential approach for wound marker analysis in *Arabidopsis thaliana*. *Journal of Chromatography B*, 871(2), 261-270.
- Grassi, D., Ferri, L., Desideri, G., Di Giosia, P., Cheli, P., Del Pinto, R., & Ferri, C. (2013). Chronic hyperuricemia, uric acid deposit and cardiovascular risk. *Current Pharmaceutical Design*, 19(13), 2432-2438.
- Guidelines Committee. (2008). *Clinical practice guidelines: Management of Gout*. Putrajaya: Ministry of Health, Malaysian Society of Rheumatology, Academy of Medicine, Malaysia.
- Hafez, R. M., Abdel-Rahman, T. M., & Naguib, R. M. (2017). Uric acid in plants and microorganisms: Biological applications and genetics-A review. *Journal of Advanced Research*, 8(5), 475-486.
- Haidari, F., Rashidi, M. R., Eshraghian, M. R., Mahboob, S. A., Shahi, M. M., & Keshavarz, S. A. (2008a). Hypouricemic and antioxidant activities of *Allium cepa* Liliaceae and quercetin in normal and hyperuricemic rats. *Saudi Medical Journal*, 29(11), 1573-1579.
- Haidari, F., Rashidi, M. R., Keshavarz, S. A., Mahboob, S. A., Eshraghian, M. R., & Shahi, M. M. (2008b). Effects of onion on serum uric acid levels and hepatic xanthine dehydrogenase/xanthine oxidase activities in hyperuricemic rats. *Pakistan Journal of Biological Sciences*, 11(14), 1779-1784.



- Haidari, F., Keshavarz, S. A., Shahi, M. M., Mahboob, S. A., & Rashidi, M. R. (2011). Effects of parsley (*Petroselinum crispum*) and its flavonol constituents, kaempferol and quercetin, on serum uric acid levels, biomarkers of oxidative stress and liver xanthine oxidoreductase activity in oxonate-induced hyperuricemic rats. *Iranian Journal of Pharmaceutical Research*, 10(4), 811-819.
- Harizal, S. N., Mansor, S. M., Hasnan, J., Tharakan, J. K. J., & Abdullah, J. (2010). Acute toxicity study of the standardized methanolic extract of *Mitragyna speciosa* Korth in rodent. *Journal of Ethnopharmacology*, 131(2), 404-409.
- Hartung, E. F. (1954). History of the use of colchicum and related medicaments in gout: with suggestions for further research. *Annals of the Rheumatic Diseases*, 13(3), 190-200.
- Hassan, N. (2001). *Recognizing gout*. Retrieved April 30, 2017, from [http://www1.utusan.com.my/utusan/info.asp?y=2001&dt=0805&sec=Kesihatan&pg=kn\\_01.htm](http://www1.utusan.com.my/utusan/info.asp?y=2001&dt=0805&sec=Kesihatan&pg=kn_01.htm)
- Hasan, K. M. M., Tamanna, N., & Haque, M. A. (2018). Biochemical and histopathological profiling of Wistar rat treated with *Brassica napus* as a supplementary feed. *Food Science and Human Wellness*, 7(1), 77-82.
- He, X. G. (2000). On-line identification of phytochemical constituents in botanical extracts by combined high-performance liquid chromatographic–diode array detection–mass spectrometric techniques. *Journal of Chromatography A*, 880(1-2), 203-232.
- Helmenstine, A.M. (2016). *Uric acid chemical structure*. Retrieved October 14, 2016 from <http://chemistry.about.com/od/factsstructures/ig/Chemical-Structures---U/Uric-Acid.htm>
- Hendriani, R., Sukandar, E. Y., Anggadiredja, K., & Sukrasno. (2016). *In vitro* evaluation of xanthine oxidase inhibitory activity of selected medicinal plants. *International Journal of Pharmaceutical and Clinical Research*, 8(4), 235-238.
- Hilaly, J.E., Israili, Z. H., & Lyoussi, B. (2004). Acute and chronic toxicological studies of *Ajuga iva* in experimental animals. *Journal of Ethnopharmacology*, 91(1), 43-50.
- Hilmi, B. A., Asmahan, M. I., & Rosman, A. (2012). Use of newly available febuxostat in a case of chronic tophaceous gout contraindicated to allopurinol and probenecid. *Medical Journal of Malaysia*, 67(1), 125-126.

- Hirano, M., Takeda, S., Hito, S., & Shimoda, H. (2017). Luteolin-rich chrysanthemum flower extract suppresses baseline serum uric acid in Japanese subjects with mild hyperuricemia. *Integrative Molecular Medicine*, 4(2), 1-5
- Hofer, D., Schwach, G., Ghaffari Tabrizi-Wizsy, N., Sadjak, A., Sturm, S., Stuppner, H., & Pfragner, R. (2013). *Christia vespertilionis* plant extracts as novel antiproliferative agent against human neuroendocrine tumor cells. *Oncology Reports*, 29(6), 2219-2226.
- Horiuchi, H., Ota, M., Nishimura, S. I., Kaneko, H., Kasahara, Y., Ohta, T., & Komoriya, K. (2000). Allopurinol induces renal toxicity by impairing pyrimidine metabolism in mice. *Life Sciences*, 66(21), 2051-2070.
- Hossain, M. A., & Rahman, S. M. (2015). Isolation and characterisation of flavonoids from the leaves of medicinal plant *Orthosiphon stamineus*. *Arabian Journal of Chemistry*, 8(2), 218-221.
- Hou, C. W., Lee, Y. C., Hung, H. F., Fu, H. W., & Jeng, K. C. (2012). Longan seed extract reduces hyperuricemia via modulating urate transporters and suppressing xanthine oxidase activity. *The American Journal of Chinese Medicine*, 40(5), 979-991.
- Huang, L., Chen, S., & Yang, M. (2012). *Euphorbia hirta* (Feiyangcao): A review on its ethnopharmacology, phytochemistry and pharmacology. *Journal of Medicinal Plants Research*, 6(39), 5176-5185.
- Huo, L. N., Wang, W., Zhang, C. Y., Shi, H. B., Liu, Y., Liu, X. H., Guo, B. H., Zhao, D. M., & Gao, H. (2015). Bioassay-guided isolation and identification of xanthine oxidase inhibitory constituents from the leaves of *Perilla frutescens*. *Molecules*, 20(10), 17848-17859.
- Iannitti, T., Capone, S., Feder, D., & Palmieri, B. (2010). Clinical use of immunosuppressants in Duchenne muscular dystrophy. *Journal of Clinical Neuromuscular Disease*, 12(1), 1-21.
- Ibrahim, M. Y., Abdul, A. B., Ibrahim, T. A. T., Abdelwahab, S. I., Elhassan, M. M., & Syam, M. M. (2010). Evaluation of acute toxicity and the effect of single injected doses of zerumbone on the kidney and liver functions in Sprague Dawley rats. *African Journal of Biotechnology*, 9(28), 4442-4450.
- Iqbal, S., & Bhanger, M. I. (2006). Effect of season and production location on antioxidant activity of *Moringa oleifera* leaves grown in Pakistan. *Journal of food Composition and Analysis*, 19(6-7), 544-551.

- Iqbal, M., Shah, M. D., Lie, C. A., & San, C. K. (2010). *Strobilanthes crispus* attenuates renal carcinogen, iron nitrilotriacetate (Fe-NTA)-mediated oxidative damage of lipids and DNA. *Molecular and Cellular Biochemistry*, 341(1-2), 271-277.
- Ishiyama, M., Tominaga, H., Shiga, M., Sasamoto, K., Ohkura, Y., & Ueno, K. (1996). A combined assay of cell viability and *in vitro* cytotoxicity with a highly water-soluble tetrazolium salt, neutral red and crystal violet. *Biological and Pharmaceutical Bulletin*, 19(11), 1518-1520.
- Jagtap, U. B., & Bapat, V. A. (2010). Artocarpus: a review of its traditional uses, phytochemistry and pharmacology. *Journal of Ethnopharmacology*, 129(2), 142-166.
- Jakše, B., Jakše, B., Pajek, M., & Pajek, J. (2019). Uric Acid and Plant-Based Nutrition. *Nutrients*, 11(8), 1736.
- Jamal, P., Azmi, S. M., Amid, A., Salleh, H. M., & Hashim, Y. Z. (2014). Development and improvement of anti-gout property from aqueous-methanol extract of *Morinda elliptica* using central composite design. *Advances in Environmental Biology*, 8, 734-743.
- Janssens, H. J., Lucassen, P. L., Van de Laar, F. A., Janssen, M., & Van de Lisdonk, E. H. (2008). Systemic corticosteroids for acute gout. *Cochrane Database of Systematic Reviews*, 2, 1-2.
- Jentzer, J. B., Alignan, M., Vaca-Garcia, C., Rigal, L., & Vilarem, G. (2015). Response surface methodology to optimise accelerated solvent extraction of steviol glycosides from *Stevia rebaudiana* Bertoni leaves. *Food Chemistry*, 166, 561-567.
- Jia, Q., Liang, Q., & Shi, Q. (2019). Astragalin suppresses inflammatory responses and bone destruction in mice with collagen-induced arthritis and in human fibroblast-like synoviocytes. *Frontiers in Pharmacology*, 10, 94.
- Jiang, Y., You, X. Y., Fu, K. L., & Yin, W. L. (2012). Effects of extract from *Mangifera indica* leaf on monosodium urate crystal-induced gouty arthritis in rats. *Evidence-Based Complementary and Alternative Medicine*, 2012, 1-6.
- Johnson, R. J., & Rideout, B. A. (2004). Uric acid and diet—insights into the epidemic of cardiovascular disease. *New England Journal of Medicine*, 350(11), 1071-1073.

- Jothy, S. L., Zuraini, Z., & Sasidharan, S. (2011). Phytochemicals screening, DPPH free radical scavenging and xanthine oxidase inhibitory activities of *Cassia fistula* seeds extract. *Journal of Medicinal Plants Research*, 5(10), 1941-1947.
- Kahn, K., Serfozo, P., & Tipton, P.A. (1997). Identification of the true product of the urate oxidase reaction. *Journal of the American Chemical Society*, 119(23), 5435-5442.
- Kaith, B. S., Bhatia, J. K., Dhiman, J., Singla, R., Mehta, P., Yadav, V., & Bhatti, M. S. (2013). Synthesis and optimization of soy protein fiber based graft copolymer through response surface methodology for removal of oil spillage. *Polymer Bulletin*, 70(11), 3155-3169.
- Kalsi, P.S., & Jagtap, S. (2013). *Pharmaceutical, medicinal and natural product chemistry*. Oxford: Alpha Science International Ltd.
- Karimi, A., Majlesi, M., & Rafieian-Kopaei, M. (2015). Herbal versus synthetic drugs; beliefs and facts. *Journal of Nephro pharmacology*, 4(1), 27-30.
- Kasarala, G., & Tillmann, H. L. (2016). Standard liver tests. *Clinical Liver Disease*, 8(1), 13-18.
- Kasolo, J.N., Bimenya, G.S., Ojok, L., Ochieng, J., & Ogwal-okeng, J.W. (2010). Phytochemicals and uses of *Moringa oleifera* leaves in Ugandan rural communities. *Journal of Medicinal Plants Research*, 4(9), 753-757.
- Kemal, C., Louis-Flamberg, P., Krupinsky-Olsen, R., & Shorter, A.L. (1987). Reductive inactivation of soybean lipoxygenase I by catechols: A possible mechanism for regulation of lipoxygenase activity. *Biochemistry*, 26(22), 7064-7072.
- Khim, K. B. (2019). *Why more younger Singaporeans are getting gout: Too much beer and protein*. Retrieved December 18, 2019 from <https://cna.lifestyle.channelnewsasia.com/wellness/gout-why-younger-singaporeans-getting-painful-form-arthritis-11302500>
- Kim, K. Y., Schumacher, H. R., Hunsche, E., Wertheimer, A. I., & Kong, S. X. (2003). A literature review of the epidemiology and treatment of acute gout. *Clinical Therapeutics*, 25(6), 1593-1617.
- Kim, W. R., Flamm, S. L., Di Bisceglie, A. M., & Bodenheimer, H. C. (2008). Serum activity of alanine aminotransferase (ALT) as an indicator of health and disease. *Hepatology*, 47(4), 1363-1370.

- King, M. A. (2000). Detection of dead cells and measurement of cell killing by flow cytometry. *Journal of Immunological Methods*, 243(1-2), 155-166.
- Klemp, P., Stansfield, S. A., Castle, B., & Robertson, M.C. (1997). Gout is on the increase in New Zealand. *Annals of the Rheumatic Diseases*, 56, 22-26.
- Kobayashi, K., Pillai, K. S., Sakuratani, Y., Abe, T., Kamata, E., & Hayashi, M. (2008). Evaluation of statistical tools used in short-term repeated dose administration toxicity studies with rodents. *The Journal of Toxicological Sciences*, 33(1), 97-104.
- Kong, L. D., Cai, Y., Huang, W. W., Cheng, C. H., & Tan, R. X. (2000). Inhibition of xanthine oxidase by some Chinese medicinal plants used to treat gout. *Journal of Ethnopharmacology*, 73(1-2), 199-207.
- Kong, L., Zhou, J., Wen, Y., Li, J., & Cheng, C. H. (2002). Aesculin possesses potent hypouricemic action in rodents but is devoid of xanthine oxidase/dehydrogenase inhibitory activity. *Planta Medica*, 68(2), 175-178.
- Kong, L.D., Yang, C., Ge, F., Wang, H.D., & Guo, Y.S. (2004). A Chinese herbal medicine Ermiao wan reduces serum uric acid level and inhibits liver xanthine dehydrogenase and xanthine oxidase in mice. *Journal of Ethnopharmacology*, 92(2-3), 325-330.
- Koshihara, Y., Fujimoto, Y., & Inoue, H. (1988). A new 5-lipoxygenase selective inhibitor derived from *Artocarpus communis* strongly inhibits arachidonic acid-induced ear edema. *Biochemical Pharmacology*, 37(11), 2161-2165.
- Kramer, H.M., & Curhan, G. (2002). The association between gout and nephrolithiasis: the National Health and Nutrition Examination Survey III, 1988-1994. *American Journal of Kidney Diseases*, 40, 37-42.
- Kumar, S., Malhotra, R., & Kumar, D. (2010). *Euphorbia hirta*: Its chemistry, traditional and medicinal uses, and pharmacological activities. *Pharmacognosy reviews*, 4(7), 58-61.
- Kumar, S., & Pandey, A. K. (2013). Chemistry and biological activities of flavonoids: an overview. *The Scientific World Journal*, 2013, 1-16.
- Kumar, A., & Azmi, W. (2014). Phytomedicine: A novel alternative for treatment of gout. *Annals of Phytomedicine*, 3(1), 80-88.
- Kuo, C. Y., Kao, E. S., Chan, K. C., Lee, H. J., Huang, T. F., & Wang, C. J. (2012). *Hibiscus sabdariffa* L. extracts reduce serum uric acid levels in oxonate-induced rats. *Journal of Functional Foods*, 4(1), 375-381.



- Kuo, C.F., Grainge, M.J., Zhang, W., & Doherty, M. (2015). Global epidemiology of gout: prevalence, incidence and risk factors. *Nature Review Rheumatology*, 11(11), 649–662.
- Lasano, N. F., Rahmat, A., Ramli, N. S., & Bakar, M. F. A. (2018). Effect of oven and microwave drying on polyphenols content and antioxidant capacity of herbal tea from *Strobilanthes crispus* leaves. *Asian Journal of Pharmaceutical and Clinical Reserach*, 11(6), 363-368.
- Latip, N. A., & Mutalib, N. A. (2019). Synergistic Interactions between *Christia vespertilionis* leaves extract and chemotherapy drug cyclophosphamide on WRL-68 cell line. *Asian Journal of Pharmaceutical Research and Development*, 7(3), 109-113.
- Lavelli, V., Peri, C., & Rizzolo, A. (2000). Antioxidant activity of tomato products as studied by model reactions using xanthine oxidase, myeloperoxidase, and copperinduced lipid peroxidation. *Journal of Agricultural and Food Chemistry*, 48(5), 5630–5639.
- Lee, J.C., Kim, H.R., Kim, J., & Jang, Y.S. (2002). Antioxidant activity of ethanol extract of the stem of *Opuntia ficus-indica* var. saboten. *Journal of Agricultural and Food Chemistry*, 50(22), 1619-1624.
- Lee, Y. W., Kim, J. D., Zheng, J., & Row, K. H. (2007). Comparisons of isoflavones from Korean and Chinese soybean and processed products. *Biochemical Engineering Journal*, 36(1), 49-53.
- Leong, C. N. A., Tako, M., Hanashiro, I., & Tamaki, H. (2008). Antioxidant flavonoid glycosides from the leaves of *Ficus pumila* L. *Food Chemistry*, 109(2), 415-420.
- Li, F., Liang, D., Yang, Z., Wang, T., Wang, W., Song, X., Guo, M., Zhou, E., Li, D., Cao, Y., & Zhang, N. (2013). Astragalín suppresses inflammatory responses via down-regulation of NF- $\kappa$ B signaling pathway in lipopolysaccharide-induced mastitis in a murine model. *International Immunopharmacology*, 17(2), 478-482.
- Li, C., Eom, T., & Jeong, Y. (2015). *Glycyrrhiza glabra* L. extract inhibits LPS-induced inflammation in RAW macrophages. *Journal of Nutritional Science and Vitaminology*, 61(5), 375-381.

- Lim, K. T., Lim, V., & Chin, J. H. (2012). Subacute oral toxicity study of ethanolic leaves extracts of *Strobilanthes crispus* in rats. *Asian Pacific Journal of Tropical Biomedicine*, 2(12), 948-952.
- Lima, R. D. C. L., Ferrari, F. C., de Souza, M. R., de Sá Pereira, B. M., de Paula, C. A., & Saúde-Guimarães, D. A. (2015). Effects of extracts of leaves from *Sparattosperma leucanthum* on hyperuricemia and gouty arthritis. *Journal of Ethnopharmacology*, 161, 194-199.
- Lin, C.N. & Shieh, W.L. (1991). Prenylflavonoids and a pyranodihydrobenzoxanthone from *Artocarpus communis*. *Phytochemistry*, 30(5), 1669–1671.
- Lin, C. N., Lu, C. M., Lin, H. C., Fang, S. C., Shieh, B. J., Hsu, M. F., Wang, J. P., Ko, F. N., & Teng, C. M. (1996). Novel antiplatelet constituents from Formosan Moraceous plants. *Journal of Natural Products*, 59(9), 834-838.
- Lin, K.C., Lin, H.Y., & Chou, P. (2000a). The interaction between uric acid level and other risk factors on the development of gout among asymptomatic hyperuricemic men in a prospective study. *Journal of Rheumatology*, 27(6), 1501 – 1505.
- Lin, K.C., Lin, H.Y., & Chou, P. (2000b). Community based epidemiological study on hyperuricemia and gout in Kin-Hu, Kinmen. *The Journal of Rheumatology*, 27(4), 1045-1050.
- Lin, C. M., Chen, C. S., Chen, C. T., Liang, Y. C., & Lin, J. K. (2002). Molecular modeling of flavonoids that inhibits xanthine oxidase. *Biochemical and Biophysical Research Communications*, 294(1), 167-172.
- Lin, Y., Xu, W., Huang, M., Xu, W., Li, H., Ye, M., Zhang, X., & Chu, K. (2015). Qualitative and quantitative analysis of phenolic acids, flavonoids and iridoid glycosides in Yinhua Kanggan tablet by UPLC-QqQ MS/MS. *Molecules*, 20(7), 12209-12228.
- Ling, X., & Bochu, W. (2014). A review of phytotherapy of gout: perspective of new pharmacological treatments. *Die Pharmazie-An International Journal of Pharmaceutical Sciences*, 69(4), 243-256.
- Ling, A. , Chin, J. , Koh, R. , Wong, Y. (2016). Investigating prostaglandin E2 and intracellular oxidative stress levels in lipopolysaccharide-stimulated RAW 264.7 macrophages upon treatment with *Strobilanthes crispus*. *International Journal of Medical and Health Sciences*, 10(5), 248 - 253.

- Liu, W., Yu, Y., Yang, R., Wan, C., Xu, B., & Cao, S. (2010). Optimization of total flavonoid compound extraction from *Gynura medica* leaf using response surface methodology and chemical composition analysis. *International Journal of Molecular Sciences*, 11(11), 4750-4763.
- Liu, L. M., Cheng, S. F., Shieh, P. C., Lee, J. C., Chen, J. J., Ho, C. T., Kuo, S. C., Kuo, D. H., Huang, L. J., & Way, T. D. (2014). The methanol extract of *Euonymus laxiflorus*, *Rubia lanceolata* and *Gardenia jasminoides* inhibits xanthine oxidase and reduce serum uric acid level in rats. *Food and Chemical Toxicology*, 70, 179-184.
- Liu, X., Huang, S., Xu, W., Zhou, A., Li, H., Zhang, R., Liu, Y., Yang, Y., & Jia, H. (2018). Association of dietary patterns and hyperuricemia: a cross-sectional study of the Yi ethnic group in China. *Food & Nutrition Research*, 62, 1380.
- Liza, M. S., Rahman, R. A., Mandana, B., Jinap, S., Rahmat, A., Zaidul, I. S. M., & Hamid, A. (2010). Supercritical carbon dioxide extraction of bioactive flavonoid from *Strobilanthes crispus* (pecah kaca). *Food and Bioprocess Technology*, 88(2-3), 319-326.
- Liza, M. S., Abdul Rahman, R., Mandana, B., Jinap, S., Rahmat, A., Zaidul, I. S. M., & Hamid, A. (2012). Supercritical fluid extraction of bioactive flavonoid from *Strobilanthes crispus* (pecah kaca) and its comparison with solvent extraction. *International Food Research Journal*, 19(2), 503-508.
- Loh, D. S. Y., Er, H. M., & Chen, Y. S. (2009). Mutagenic and antimutagenic activities of aqueous and methanol extracts of *Euphorbia hirta*. *Journal of Ethnopharmacology*, 126(3), 406-414.
- Luk, A. J., & Simkin, P. A. (2005). Epidemiology of hyperuricemia and gout. *The American Journal of Managed Care*, 11(15), S435-S442.
- Lundstedt, T., Seifert, E., Abramo, L., Thelin, B., Nyström, Å., Pettersen, J., & Bergman, R. (1998). Experimental design and optimization. *Chemometrics and Intelligent Laboratory Systems*, 42(1), 3-40.
- Lyu, L. C., Hsu, C. Y., Yeh, C. Y., Lee, M. S., Huang, S. H., & Chen, C. L. (2003). A case-control study of the association of diet and obesity with gout in Taiwan. *The American journal of clinical nutrition*, 78(4), 690-701.
- Ma, F., Zhao, Y., Gong, X., Xie, Y., & Zhou, X. (2014). Optimization of quercitrin and total flavonoids extraction from *Herba Polygoni Capitati* by response surface methodology. *Pharmacognosy Magazine*, 10(1), S57.



- Majdan, M., & Borys, O. (2010). Dna i schorzenia towarzyszące podwyższonemu stężeniu kwasu moczowego. *Rocz. Pomor. Academic Medicine*, 56, 34-39.
- Majeed, M., Hussain, A. I., Chatha, S. A., Khosa, M. K., Kamal, G. M., Kamal, M. A., Zhang, X., & Liu, M. (2016). Optimization protocol for the extraction of antioxidant components from *Origanum vulgare* leaves using response surface methodology. *Saudi Journal of Biological Sciences*, 23(3), 389-396.
- Major, T. J., Topless, R. K., Dalbeth, N., & Merriman, T. R. (2018). Evaluation of the diet wide contribution to serum urate levels: meta-analysis of population based cohorts. *BMJ*, 363, k3951.
- Mandell, B. F. (2008). Clinical manifestations of hyperuricemia and gout. *Cleveland Clinic Journal of Medicine*, 75(5), S5-S8.
- Martinon, F., Pétrilli, V., Mayor, A., Tardivel, A., & Tschopp, J. (2006). Gout-associated uric acid crystals activate the NALP3 inflammasome. *Nature*, 440, 237-241.
- Matsuo, H., Chiba, T., Nagamori, S., Nakayama, A., Domoto, H., Phetdee, K., Wiriyaermskul, P., Kikuchi, Y., Oda, T., Nishiyama, J., Nakamura, T., Morimoto, Y., Kamakura, K., Sakurai, Y., Nonoyama, S., Kanai, Y., & Shinomiya, N. (2008). Mutations in glucose transporter 9 gene SLC2A9 cause renal hypouricemia. *The American Journal of Human Genetics*, 83(6), 744-751.
- Mat-Salleh, K., & Latiff, A. (2005). *Tumbuhan Ubatan Malaysia*. Malaysia: Universiti Kebangsaan Malaysia, 254.
- Mbikay, M. (2012). Therapeutic potential of *Moringa oleifera* leaves in chronic hyperglycemia and dyslipidemia: a review. *Frontiers in Pharmacology*, 3, 1–12.
- McAdams DeMarco, M. A., Maynard, J. W., Huizinga, M. M., Baer, A. N., Köttgen, A., Gelber, A. C., & Coresh, J. (2011). Obesity and younger age at gout onset in a community-based cohort. *Arthritis Care & Research*, 63(8), 1108-1114.
- McLean, S., Reynolds, W. F., Tinto, W. F., Chan, W. R., & Shepherd, V. (1996). Complete <sup>13</sup>C and <sup>1</sup>H spectral assignments of prenylated flavonoids and a hydroxy fatty acid from the leaves of Caribbean *Artocarpus communis*. *Magnetic Resonance in Chemistry*, 34(9), 719-722.
- Media, F. G. (2017). *Gout: More common in men than in women?*. Retrieved December 18, 2019 from <https://voiz.asia/en/15597>

- Mehmood, A., Zhao, L., Ishaq, M., Safdar, B., Wang, C., & Nadeem, M. (2018). Optimization of total phenolic contents, antioxidant, and *in-vitro* xanthine oxidase inhibitory activity of sunflower head. *CyTA-Journal of Food*, 16(1), 957-964.
- Meng, Z. Q., Tang, Z. H., Yan, Y. X., Guo, C. R., Cao, L., Ding, G., Huang, W. Z., Wang, Z. Z., Wang, K. D. G., Xiao, W., & Yang, Z. L. (2014). Study on the anti-gout activity of chlorogenic acid: Improvement on hyperuricemia and gouty inflammation. *The American Journal of Chinese Medicine*, 42(6), 1471-1483.
- Mirghani, M. E. S., Liyana, Y., & Parveen, J. (2012). Bioactivity analysis of lemongrass (*Cymbopogon citratus*) essential oil. *International Food Research Journal*, 19(2), 569-575.
- Mittag, F., & Wuenschel, M. (2011). Giant gouty tophi of the hand and wrist. *Orthopedics*, 34(11), 790-792.
- Mo, S. F., Zhou, F., Lv, Y. Z., Hu, Q. H., Zhang, D. M., & Kong, L. D. (2007). Hypouricemic action of selected flavonoids in mice: structure-activity relationships. *Biological and Pharmaceutical Bulletin*, 30(8), 1551-1556.
- Mohamed, D.A., & Al-Okbi, S. Y. (2008). Evaluation of anti-gout activity of some plant food extracts. *Polish Journal of Food and Nutrition Science*, 58(3), 389-395.
- Mohamed, E. A. H., Siddiqui, M. J. A., Ang, L. F., Sadikun, A., Chan, S. H., Tan, S. C., Asmawi, M. Z., & Yam, M. F. (2012). Potent  $\alpha$ -glucosidase and  $\alpha$ -amylase inhibitory activities of standardized 50% ethanolic extracts and sinensetin from *Orthosiphon stamineus* Benth as anti-diabetic mechanism. *BMC Complementary and Alternative Medicine*, 12(1), 176.
- Mohd, A., Das Gupta, E., Loh, Y. L., Gandhi, C., D'Souza, B., & Gun, S. C. (2011). Clinical characteristics of gout: a hospital case series. *Malaysian Family Physician: The Official Journal of The Academy of Family Physicians of Malaysia*, 6(2-3), 72.
- Molad, Y. (2002). Update on colchicine and its mechanism of action. *Current Rheumatology Reports*, 4(3), 252-256.
- Montgomery, D.C. (1997). *Design and Analysis of Experiments*, New York: Wiley.

- Mosmann, T. (1983). Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *Journal of Immunological Methods*, 65(1-2), 55-63.
- Mujumdar, A.M., & Misar, A. V. (2004). Anti-inflammatory activity of *Jatropha curcas* roots in mice and rats. *Journal of Ethnopharmacology*, 90(1), 11-15.
- Murata, K., Nakao, K., Hirata, N., Namba, K., Nomi, T., Kitamura, Y., Moriyama, K., Shintani, T., Linuma, M., & Matsuda, H. (2009). Hydroxychavicol: a potent xanthine oxidase inhibitor obtained from the leaves of betel, *Piper betle*. *Journal of Natural Medicines*, 63(3), 355-359.
- Murugaiyah, V., & Chan, K. L. (2006). Antihyperuricemic lignans from the leaves of *Phyllanthus niruri*. *Planta Medica*, 72(14), 1262-1267.
- Murugaiyah, V., & Chan, K. L. (2009). Mechanisms of antihyperuricemic effect of *Phyllanthus niruri* and its lignan constituents. *Journal of Ethnopharmacology*, 124(2), 233-239.
- Mutee, A. F., Salhimi, S. M., Yam, M. F., Lim, C. P., Abdullah, G. Z., Ameer, O. Z., Abdul Karim, M. F., & Asmawi, M.Z. (2010). *In vivo* anti-inflammatory and *in vitro* antioxidant activities of *Peperomia pellucida*. *International Journal of Pharmacology*, 6(5), 686-690.
- Nana, F. W., Hilou, A., Millogo, J. F., & Nacoulma, O. G. (2012). Phytochemical composition, antioxidant and xanthine oxidase inhibitory activities of *Amaranthus cruentus* L. and *Amaranthus hybridus* L. extracts. *Pharmaceuticals*, 5(6), 613-628.
- Naseem, S.A., Muhammad, F., Muzammil, H.N., Kouser, B.M., & Aurangzeb, H. (2006). Activity of polyphenolic plant extracts as scavengers of free radicals and inhibitors of xanthine oxidase. *The Journal of Basic and Applied Sciences*, 2(1), 1-6.
- Nessa, F., Ismail, Z., & Mohamed, N. (2010). Xanthine oxidase inhibitory activities of extracts and flavonoids of the leaves of *Blumea balsamifera*. *Pharmaceutical Biology*, 48(12), 1405-1412.
- Neto, B. B., Scarminio, I. S., & Bruns, R. E. (1996). *Planejamento e otimização de experimentos*. Ed. da UNICAMP.
- Nguyen, M.T.T., Awale, S., Tezuka, Y., Tran, Q.L., Watanabe, H., & Kadota, S. (2004). Xanthine oxidase inhibitory activity of Vietnamese medicinal plants. *Biological and Pharmaceutical Bulletin*, 27(9), 1414-1421.

- Nguyen, M. K. H., Nguyen, H. X., Nguyen, N. T., & Nguyen, M. T. T. (2014). Total Polyphenol contents of An Giang medicinal plants. *Science and Technology Development Journal*, 17(2), 5-9.
- Nguyen, T. D., Thuong, P. T., Hwang, I. H., Hoang, T. K. H., Nguyen, M. K., Nguyen, H. A., & Na, M. (2017). Anti-hyperuricemic, anti-inflammatory and analgesic effects of *Siegesbeckia orientalis* L. resulting from the fraction with high phenolic content. *BMC Complementary and Alternative Medicine*, 17(1), 191.
- Nguyen, T. T., Mai, H. T. T., Nguyen, T. N., Tran, Q. D., & Tran, T. D. (2019). Optimization of extraction of total phenolic and total flavonoid from seeds of *Alpinia blepharocalyx* K. Schum. using the response surface methodology. *Vietnam Journal of Science and Technology*, 57(3B), 105-115.
- Nguyen-Pouppln, J., Tran, H., Tran, H., Phan, T. A., Dolecek, C., Farrar, J., Tran, T.H., Caron, P., Bodo, B., & Grellier, P. (2007). Antimalarial and cytotoxic activities of ethnopharmacologically selected medicinal plants from South Vietnam. *Journal of Ethnopharmacology*, 109(3), 417-427.
- Nile, S. H., & Park, S. W. (2014). HPTLC analysis, antioxidant and antigout activity of Indian plants. *Iranian Journal of Pharmaceutical Research*, 13(2), 531-539.
- Nisar, B., Sultan, A., & Rubab, S. L. (2017). Comparison of medicinally important natural products versus synthetic drugs-a short commentary. *Natural Products Chemistry Research*, 6(2), 1-2.
- Noraida, A. (2005). *Penyembuhan Semula Jadi dengan Herba*. Kuala Lumpur : PTS Millennia Sdn. Bhd.
- Nurraihana, H., & Norfarizan-Hanoon, N. A. (2013). Phytochemistry, pharmacology and toxicology properties of *Strobilanthes crispus*. *International Food Research Journal*, 20(5), 2045-2056.
- Nurul, S. A. S., Hazilawati, H., Mohd, R. S., Mohd, F. H. R., Noordin, M. M., & Norhaizan, M. E. (2018). Subacute oral toxicity assesment of ethanol extract of *Mariposa christia vespertilionis* leaves in male Sprague Dawley rats. *Toxicological Research*, 34(2), 85-95.
- OECD. (2002). *Guidelines for the testing of chemicals / section 4: health effects test no. 423: acute oral toxicity - acute toxic class method*. Paris, France: Organization for Economic Cooperation and Development.

- Ogbole, O. O., Segun, P. A., & Adeniji, A. J. (2017). *In vitro* cytotoxic activity of medicinal plants from Nigeria ethnomedicine on Rhabdomyosarcoma cancer cell line and HPLC analysis of active extracts. *BMC Complementary and Alternative Medicine*, 17(1), 494.
- Olowa, L., & Demayo, C. G. (2015). Ethnobotanical uses of medicinal plants among the Muslim Maranaos in Iligan City, Mindanao, Philippines. *Advances in Environmental Biology*, 9(27), 204-216.
- Oskoueian, E., Abdullah, N., Hendra, R., & Karimi, E. (2011). Bioactive compounds, antioxidant, xanthine oxidase inhibitory, tyrosinase inhibitory and anti-inflammatory activities of selected agro-industrial by-products. *International Journal of Molecular Sciences*, 12(12), 8610-8625.
- Osman, N. I., Sidik, N. J., Awal, A., Adam, N. A. M., & Rezali, N. I. (2016). *In vitro* xanthine oxidase and albumin denaturation inhibition assay of *Barringtonia racemosa* L. and total phenolic content analysis for potential anti-inflammatory use in gouty arthritis. *Journal of Intercultural Ethnopharmacology*, 5(4), 343-349.
- Osman, M. S., Ghani, Z. A., Ismail, N. F., Razak, N. A. A., Jaapar, J., & Ariff, M. A. M. (2017). Qualitative comparison of active compounds between red and green *Mariposa christia vespertillonis* leaves extracts. *AIP Conference Proceedings*, 1885(1), 020282.
- Owen, P. L., & Johns, T. (1999). Xanthine oxidase inhibitory activity of northeastern North American plant remedies used for gout. *Journal of Ethnopharmacology*, 64(2), 149-160.
- Pacher, P., Nivorozhkin, A., & Szabo, C. (2006). Therapeutic effect of xanthine oxidase inhibitors: Renaissance half a century after the discovery of allopurinol. *Pharmacology Review*, 58(1), 87-114.
- Palu, A., Deng, S., West, B., & Jensen, J. (2009). Xanthine oxidase inhibiting effects of noni (*Morinda citrifolia*) fruit juice. *Phytotherapy Research*, 23(12), 1790-1791.
- Park, J. E., Yeom, Z., Park, K. T., Han, E. H., Yu, H. J., Kang, H. S., & Lim, Y. H. (2018). Hypouricemic effect of ethanol extract of *Aster glehni* leaves in potassium oxonate-induced hyperuricemic rats. *Clinical Nutrition Research*, 7(2), 126-135.
- Parrotta, J. A. (2001). *Healing plants of peninsular India*. UK: CABI publishing.



- Pascual, E., Addadi, L., Andrés, M., & Sivera, F. (2015). Mechanisms of crystal formation in gout—a structural approach. *Nature Reviews Rheumatology*, 11(12), 725-730.
- Paul, B. J., & James, R. (2017). Gout: an Asia-Pacific update. *International Journal of Rheumatic Diseases*, 20(4), 407-416.
- Peñarrieta, J. M., Alvarado, J. A., Bergenståhl, B., & Åkesson, B. (2007). Spectrophotometric methods for the measurement of total phenolic compounds and total flavonoids in foods. *Revista Boliviana de Química*, 24(1), 5-9.
- Perez, C., Pauli, A., & Bazerque, P. (1990). Antibiotic assay by agar-well diffusion method. *Acta Biologiae Medicine Experimentalis*, 15, 113-115.
- Perez-Ruiz, F., Alonso-Ruiz, A., Calabozo, M., Herrero-Beites, A., Garcia-Erauskin, G., & Ruiz-Lucea, E. (1998). Efficacy of allopurinol and benzbromarone for the control of hyperuricaemia. A pathogenic approach to the treatment of primary chronic gout. *Annals of The Rheumatic Diseases*, 57(9), 545-549.
- Perez-Ruiz, F., Mielgo, F. J. G., & Herrero-Beites, A. M. (2000). Optimisation of the treatment of acute gout. *BioDrugs*, 13(6), 415-423.
- Ping, K. Y., Darah, I., Chen, Y., Sreeramanan, S., & Sasidharan, S. (2013). Acute and subchronic toxicity study of *Euphorbia hirta* L. methanol extract in rats. *BioMed Research International*, 2013, 1-14.
- Poór, G., & Mituszova, M. (2003). History, classification and epidemiology of crystal-related arthropathies. *Rheumatology*, 2, 1893-1902.
- Pradhan, C., Mohanty, M., & Rout, A. (2012). Phytochemical screening and comparative bioefficacy assessment of *Artocarpus altilis* leaf extracts for antimicrobial activity. *Frontiers in Life Science*, 6(3-4), 71-76.
- Prasad, K. N., Hassan, F. A., Yang, B., Kong, K. W., Ramanan, R. N., Azlan, A., & Ismail, A. (2011). Response surface optimisation for the extraction of phenolic compounds and antioxidant capacities of underutilised *Mangifera pajang* Kosterm. peels. *Food Chemistry*, 128(4), 1121-1127.
- Punzi, L., & So, A. (2013). Serum uric acid and gout: from the past to molecular biology. *Current Medical Research and Opinion*, 29(3), 3-8.
- Punzi, L., Scanu, A., Spinella, P., Galozzi, P., & Oliviero, F. (2019). One year in review 2018: gout. *Clinical and Experimental Rheumatology*, 37(1), 1-11.
- Qurie, A., & Musa, R. (2018). *Allopurinol*. Treasure Island: StatPearls Publishing.

- Rackova, L., Oblozinsky, M., Kostalova, D., Kettmann, V., & Bezakova, L. (2007). Free radical scavenging activity and lipoxygenase inhibition of *Mahonia aquifolium* extract and isoquinoline alkaloids. *Journal of Inflammation*, 4, 15-21.
- Radojkovića, M., Zekovića, Z., Jokićb, S. & Vidovića, S. (2012). Determination of optimal extraction parameters of mulberry leaves using response surface methodology (RSM). *Romanian Biotechnological Letters*, 17(3), 7295-7308.
- Ragone, D. (1997). *Breadfruit (Artocarpus altilis)*. Hawai'i, USA: Permanent Agriculture Resources.
- Rahman, M. (2018). *Application of computational methods in isolation of plant secondary metabolites*. In *Computational Phytochemistry*. Elsevier, 107-139.
- Rahmat, A., Edrini, S., Akim, A.M., Ismail, P., Hin, T.Y.Y., & Abu Bakar, M.F. (2006). Anticarcinogenic properties of *Strobilanthes crispus* extracts and its compounds *in vitro*. *International Journal of Cancer Research*, 2(1), 47-49.
- Rajendran, R., & Krishnakumar, E. (2010). Anti-arthritis activity of *Premna serratifolia* Linn., wood against adjuvant induced arthritis. *Avicenna Journal of Medical Biotechnology*, 2(2), 101-106.
- Raju, R., Joseph, S., Scaria, S., Mathews, S. M., & Umamaheshwari, M. (2012). Effect of the fractions of *Erythrina stricta* leaf extract on serum urate levels and Xb/Xdh activities in oxonate-induced hyperuricaemic mice. *Journal of Applied Pharmaceutical Science*, 2(2), 89-94.
- Randhir, R., Lin, Y. T., & Shetty, K. (2004). Phenolics, their antioxidant and antimicrobial activity in dark germinated fenugreek sprouts in response to peptide and phytochemical elicitors. *Asia Pacific Journal of Clinical Nutrition*, 13(3), 295-307.
- Ranu, H., Jiang, J., & Ming, P. S. (2011). A case series of allopurinol-induced toxic epidermal necrolysis. *Indian Journal of Dermatology*, 56(1), 74-76.
- Rastogi, R. P., & Mehrotra (2002). *Compendium of Indian Medicinal Plants: 1985-1989*. New Delhi: Central Drug Research Institute, Lucknow.
- Raza, M., Al-Shabanah, O.A., El-Hadiyah, T.M., & AlMajed, A.A. (2002). Effect of prolonged vigabatrin treatment on hematological and biochemical parameters in plasma, liver and kidney of Swiss albino mice. *Scientia Pharmaceutica*, 70(2), 135-145.

- Reddy, P., & Lokesh, B. R. (1994). Studies on anti-inflammatory activity of spice principles and dietary n-3 polyunsaturated fatty acids on carrageenan-induced inflammation in rats. *Annals of Nutrition and Metabolism*, 38(6), 349-358.
- Reinders, M. K., & Brouwers, J. R. B. J. (2008). Benzbromarone: an old drug with new perspectives; update of its clinical pharmacology. In *Practice Research in The Field of Gout*. Netherlands: Ponsen & Looijen BV. 29-44.
- Reinders, M. K., Haagsma, C., Jansen, T. T. A., Van Roon, E. N., Delsing, J., Van de Laar, M. A. F. J., & Brouwers, J. R. B. J. (2009). A randomised controlled trial on the efficacy and tolerability with dose escalation of allopurinol 300–600 mg/day versus benzbromarone 100–200 mg/day in patients with gout. *Annals of the Rheumatic Diseases*, 68(6), 892-897.
- Rees, F., Hui, M., & Doherty, M. (2014). Optimizing current treatment of gout. *Nature Reviews Rheumatology*, 10(5), 271-283.
- Rezvanian, M., Amin, M. C. I. M., & Ng, S. F. (2016). Development and physicochemical characterization of alginate composite film loaded with simvastatin as a potential wound dressing. *Carbohydrate polymers*, 137, 295-304.
- Richette, P., & Bardin (2010). Gout. *The lancet*, 375(9711), 318–328.
- Riss, T. L., Moravec, R. A., Niles, A. L., Duellman, S., Benink, H. A., Worzella, T. J., & Minor, L. (2016). Cell viability assays. In *Assay Guidance Manual [Internet]*. Eli Lilly & Company and the National Center for Advancing Translational Sciences.
- Roberts, W. N., Liang, M. H., & Stern, S. H. (1987). Colchicine in acute gout: reassessment of risks and benefits. *JAMA*, 257(14), 1920-1922.
- Robinson, P.C., Merriman, T.R., Herbison, P., & Highton, J. (2013). Hospital admissions associated with gout and their comorbidities in New Zealand and England 1999-2009. *Rheumatology*, 52(1), 118-126.
- Roddy, E., & Doherty, M. (2010). Gout. Epidemiology of gout. *Arthritis Research and Therapy*, 12(6), 223.
- Roohbakhsh, A., Shamsara, J., Khayyat, M. H., & Karimi, G. (2009). Inhibition of xanthine oxidase by some Iranian plant remedies used for gout. *Pharmacologyonline*, 3, 1031-1036.



- Rozenberg, S., Lang, T., Laatar, A., Koeger, A. C., Orcel, P., & Bourgerois, P. (1996). Diversity of opinions on the management of gout in France. A survey of 750 rheumatologists. *Revue du rhumatisme*, 63(4), 255-261.
- Ruoff, G., & Edwards, N.L. (2016). Overview of serum uric acid treatment targets in gout: why less than 6 mg/dL?. *Postgraduate Medicine*, 128(7), 706-715.
- Saag, K.G., & Mikuls, T.R. (2005). Recent advances in the epidemiology of gout. *Current Rheumatology Reports*, 7, 235-241.
- Sabina, E. P., & Rasool, M. (2008). An *in vivo* and *in vitro* potential of Indian ayurvedic herbal formulation Triphala on experimental gouty arthritis in mice. *Vascular Pharmacology*, 48(1), 14-20.
- Sabina, E. P., Nagar, S., & Rasool, M. (2011). A role of piperine on monosodium urate crystal-induced inflammation—an experimental model of gouty arthritis. *Inflammation*, 34(3), 184-192.
- Samarakoon, S. R., Thabrew, I., Galhena, P. B., De Silva, D., & Tennekoon, K. H. (2010). A comparison of the cytotoxic potential of standardized aqueous and ethanolic extracts of a polyherbal mixture comprised of *Nigella sativa* (seeds), *Hemidesmus indicus* (roots) and *Smilax glabra* (rhizome). *Pharmacognosy Research*, 2(6), 335-342.
- Sang, M., Du, G., Hao, J., Wang, L., Liu, E., Zhang, Y., Wang, T., Gao, X., & Han, L. (2017). Modeling and optimizing inhibitory activities of *Nelumbinis folium* extract on xanthine oxidase using response surface methodology. *Journal of Pharmaceutical and Biomedical Analysis*, 139, 37-43.
- Sapra, S., Bhalla, Y., Sharma, S., Singh, G., Nepali, K., Budhiraja, A., & Dhar, K. L. (2013). Colchicine and its various physicochemical and biological aspects. *Medicinal Chemistry Research*, 22(2), 531-547.
- Sarvaiya, V.N., Sadariya, K. A., Pancha, P. G., Thaker, A. M., Patel, A. C., & Prajapati, A. S. (2015). Evaluation of antigout activity of *Phyllanthus emblica* fruit extracts on potassium oxonate-induced gout rat model. *Veterinary World*, 8(10), 1230-1236.
- Saw, L.G., & Chung, R.C.K. (2015). The flora of Malaysia projects. *Rodriguesia*, 66(4), 947-960.
- Schofield, P., Mbugua, D. M., & Pell, A. N. (2001). Analysis of condensed tannins: a review. *Animal Feed Science and Technology*, 91(1-2), 21-40.

- Sharma, N., Samarakoon, K., Gyawali, R., Park, Y. H., Lee, S. J., Oh, S. J., Lee, T. H., & Jeong, D. (2014). Evaluation of the antioxidant, anti-inflammatory, and anticancer activities of *Euphorbia hirta* ethanolic extract. *Molecules*, 19(9), 14567-14581.
- Sharp, P. E., La Regina, M. C., & Suckow, M. (1998). The laboratory rat CRC Press. Boca Raton (collection: *The Laboratory Animal Pocket Reference Series*), 214.
- Sheng, Z. L., Wan, P. F., Dong, C. L., & Li, Y. H. (2013). Optimization of total flavonoids content extracted from Flos Populi using response surface methodology. *Industrial Crops and Products*, 43, 778-786.
- Sherman, M. R., Saifer, M. G. P., & Ruiz, F. P. (2008). PEG-uricase in the management of treatment-resistant gout and hyperuricemia. *Advanced Drug Delivery Reviews*, 60(1), 59-68.
- Sheu, S.Y., Fu, Y. T., Huang, W. D., Chen, Y. A., Lei, Y. C., Yao, C. H., Hsu, F. L., & Kuo, T. F. (2016). Evaluation of xanthine oxidase inhibitory potential and *in vivo* hypouricemic activity of *Dimocarpus longan* lour. extracts. *Pharmacognosy Magazine*, 12, 206-212.
- Shi, J., Yu, J., Pohorly, J., Young, J. C., Bryan, M., & Wu, Y. (2003). Optimization of the extraction of polyphenols from grape seed meal by aqueous ethanol solution. *Food, Agriculture & Environment*, 1(2), 42-47.
- Shi, Y. C., Lin, K. S., Jhai, Y. F., Lee, B. H., Han, Y., Cui, Z., Hsu, W. H., & Wu, S.C. (2016). Miracle fruit (*Synsepalum dulcificum*) exhibits as a novel anti-hyperuricaemia agent. *Molecules*, 21(2), 140-150.
- Shuid, A. N., Siang, L. K., Chin, T. G., Muhammad, N., Mohamed, N., & Soelaiman, I. N. (2011). Acute and subacute toxicity studies of *Eurycoma longifolia* in male rats. *International Journal of Pharmacology*, 7(5), 641-646.
- Shukor, N. A. A., Ablat, A., Muhamad, N. A., & Mohamad, J. (2018). *In vitro* antioxidant and *in vivo* xanthine oxidase inhibitory activities of *Pandanus amaryllifolius* in potassium oxonate-induced hyperuricemic rats. *International Journal of Food Science & Technology*, 53(6), 1476-1485.
- Sikarwar, M. S., Hui, B. J., Subramaniam, K., Valeisamy, B. D., Yean, L. K., & Balaji, K. (2014). Antioxidant activity of *Artocarpus altilis* (Parkinson) Fosberg leaves. *Free Radicals & Antioxidants*, 4(2), 33-39.

- Silva, E. M., Rogez, H., & Larondelle, Y. (2007). Optimization of extraction of phenolics from *Inga edulis* leaves using response surface methodology. *Separation and Purification Technology*, 55(3), 381-387.
- Singh, B. N., Singh, B. R., Singh, R. L., Prakash, D., Dhakarey, R., Upadhyay, G., & Singh, H. B. (2009). Oxidative DNA damage protective activity, antioxidant and anti-quorum sensing potentials of *Moringa oleifera*. *Food and Chemical Toxicology*, 47(6), 1109-1116.
- Singh, J. A. (2013). Racial and gender disparities among patients with gout. *Current Rheumatology Reports*, 15(2), 307.
- Singh, G., & Kumar, P. (2013). Phytochemical study and screening for antimicrobial activity of flavonoids of *Euphorbia hirta*. *International Journal of Applied and Basic Medical Research*, 3(2), 111-116.
- Singh, M. K., Gidwani, B., Gupta, A., Dhongade, H., Kashyap, P. P., & Tripathi, D. K. (2015). A review of the medicinal plants of genus *Orthosiphon* (Lamiaceae). *International Journal of Biological Chemistry*, 9(6), 318-331.
- Siu, Y. P., Leung, K. T., Tong, M. K. H., & Kwan, T. H. (2006). Use of allopurinol in slowing the progression of renal disease through its ability to lower serum uric acid level. *American Journal of Kidney Diseases*, 47(1), 51-59.
- Siva, C., Velazquez, C., Mody, A., & Brasington, R. (2003). Diagnosing acute monoarthritis in adults: a practical approach for the family physician. *American Family Physician*, 68(1), 83-92.
- Skibo, E. B. (1986). Noncompetitive and irreversible inhibition of xanthine oxidase by benzimidazole analogs acting at the functional flavin adenine dinucleotide cofactor. *Biochemistry*, 25(15), 4189-4194.
- Smith, E. U. R., Diaz-Torne, C., Perez-Ruiz, F., & March, L. M. (2010). Epidemiology of gout: an update. *Best Practice & Research Clinical Rheumatology*, 24(6), 811-827.
- Soediro, I., Pellcuer, J., Andary, C., & Privat, G. (1983). *Strobilanthes crispus*, (L) BL. I. Isolation and Identification of the caffeic acid derivative, verbascoside, 99: 50302q. *Acta Pharmaceutica Indonesia*, 8, 1-10.
- Soediro, I., Pellecuer, J., Andary, C., & Privat, G. (1987). *Strobilanthes crispus* (L) Bl I: Pemeriksaan asam fenolat. *Acta Pharmaceutica Indonesia*, 12, 1-7.

- Somchit, M. N., Mak, J. H., Bustamam, A. A., Zuraini, A., Arifah, A. K., Adam, Y., & Zakaria, Z.A. (2012). Zerumbone isolated from *Zingiber zerumbet* inhibits inflammation and pain in rats. *Journal of Medicinal Plants Research*, 6(2), 177-180.
- Sood, S. K., Bhardwaj, R., & Lakhanpal, T. N. (2005). *Ethnic Indian plants in cure of diabetes*. India: Scientific publishers.
- Soonthornsit, N., Pitaksutheepong, C., Hemstapat, W., Utaisinchaoen, P., & Pitaksuteepong, T. (2017). *In vitro* anti-inflammatory activity of *Morus alba* L. stem extract in LPS-stimulated RAW 264.7 cells. *Evidence-Based Complementary and Alternative Medicine*, 2017, 1-8.
- Soromou, L. W., Chen, N., Jiang, L., Huo, M., Wei, M., Chu, X., Millimouno, F. M., Feng, H., Sidime, Y., & Deng, X. (2012). Astragalin attenuates lipopolysaccharide-induced inflammatory responses by down-regulating NF- $\kappa$ B signaling pathway. *Biochemical and Biophysical Research Communications*, 419(2), 256-261.
- Souza, A. S., dos Santos, W. N., & Ferreira, S. L. (2005). Application of Box–Behnken design in the optimisation of an on-line pre-concentration system using knotted reactor for cadmium determination by flame atomic absorption spectrometry. *Spectrochimica Acta Part B: Atomic Spectroscopy*, 60(5), 737-742.
- Sowndhararajan, K., Joseph, J.M., & Rajendrakumaran, D. (2012). *In vitro* xanthine oxidase inhibitory activity of methanol extracts of *Erythrina indica* Lam. leaves and stem bark. *Asian Pacific Journal of Tropical Biomedicine*, 2(3), 1415-S1417.
- Spanou, C., Veskoukis, A. S., Kerasioti, T., Kontou, M., Angelis, A., Aligiannis, N., Skaltsounis, A. L., & Kouretas, D. (2012). Flavonoid glycosides isolated from unique legume plant extracts as novel inhibitors of xanthine oxidase. *PLoS One*, 7(3), e32214.
- Sreelatha, S., & Padma, P. R. (2009). Antioxidant activity and total phenolic content of *Moringa oleifera* leaves in two stages of maturity. *Plant Foods for Human Nutrition*, 64(4), 303-311.
- Sreeramulu, D., & Raghunath, M. (2010). Antioxidant activity and phenolic content of roots, tubers and vegetables commonly consumed in India. *Food Research International*, 43(4), 1017-1020.

- Sriplang, K., Adisakwattana, S., Rungsipipat, A., & Yibchok-Anun, S. (2007). Effects of *Orthosiphon stamineus* aqueous extract on plasma glucose concentration and lipid profile in normal and streptozotocin-induced diabetic rats. *Journal of Ethnopharmacology*, 109(3), 510-514.
- Sri-Wahjuni, Putra-Manuaba, I. B., Rahayu-Artini, N. P., & Wahyu-Dwijani, S. (2012). Uric acid inhibition activity of *Annona muricata* L. leave extract in hyperuricemia induced wistar rat. *Advanced Pure and Applied Chemistry*, 2(1), 86-90.
- Steffensen, S. K., Rinnan, Å., Mortensen, A. G., Laursen, B., de Troiani, R. M., Noellemeyer, E. J., Janovska, D., Dusek, K., Délano-Frier, J., Taberner, A., Christophersen, C., & Fomsgaard, I. S. (2011). Variations in the polyphenol content of seeds of field grown *Amaranthus* genotypes. *Food Chemistry*, 129(1), 131-138.
- Stone, V., Johnston, H., & Schins, R. P. (2009). Development of *in vitro* systems for nanotoxicology: methodological considerations. *Critical Reviews in Toxicology*, 39(7), 613-626.
- Strazzullo, P., & Puig, J.G. (2007). Uric acid and oxidative stress: relative impact on cardiovascular risk. *Nutrition, Metabolism and Cardiovascular Diseases*, 17, 409-414.
- Sulaiman, W., Zuki, N. W. M., Zamri, N., Suganthan, S., Abdullah, A. C., & Seung, O. P. (2019). Epidemiology and management of gout patients attending Rheumatology Tertiary Centre in Perak, Malaysia. *Asian Journal of Medicine and Health Sciences*, 2(1), 20-27.
- Sunarni, T., Leviana, F., Fidrianny, I., Iwo, M. I., & Wirasutisna, K. R. (2015). Antihyperuricemic activity of four plants Annonaceae using hyperuricemic rats model and enzyme assay. *Asian Journal of Pharmaceutical and Clinical Research*, 8(6), 250-253.
- Surveswaran, S., Cai, Y.Z., Corke, H., & Sun, M. (2007). Systematic evaluation of natural phenolic antioxidants from 133 Indian medicinal plants. *Food Chemistry*, 102, 938-953.
- Sushmita, & Nayeem, N. (2013). *Artocarpus altilis*: Over view of a plant which is referred to as bread fruit. *International Journal Pharmaceutical Sciences Letters*, 3(5), 273-276.



- Sutaria, S., Katbamna, R., & Underwood, M. (2006). Effectiveness of interventions for the treatment of acute and prevention of recurrent gout—a systematic review. *Rheumatology*, 45(11), 1422-1431.
- Taciak, B., Białasek, M., Braniewska, A., Sas, Z., Sawicka, P., Kiraga, Ł., Rygiel, T., & Król, M. (2018). Evaluation of phenotypic and functional stability of RAW 264.7 cell line through serial passages. *PloS one*, 13(6), e0198943.
- Tan, M. J., Ye, J. M., Turner, N., Hohnen-Behrens, C., Ke, C. Q., Tang, C. P., Chen, T., Weiss, H. C., Gesing, E. R., Rowland, A., James, D. E., & Ye, Y. (2008). Antidiabetic activities of triterpenoids isolated from bitter melon associated with activation of the AMPK pathway. *Chemistry & Biology*, 15(3), 263-273.
- Tapsell, L. C., Hemphill, I., Cobiac, L., Patch, C. S., Sullivan, D. R., Fenech, M., Patch, C. S., Roodenrys, S., Keogh, J. B., Clifton, P. M., Williams, P. G., Fazio, V. A., & Inge, K. E. (2006). Health benefits of herbs and spices: The past, the present, the future. *The Medical Journal of Australia*, 185(4), S4–S24.
- Tausche, A. K., Jansen, T. L., Schröder, H. E., Bornstein, S. R., Aringer, M., & Müller-Ladner, U. (2009). Gout—current diagnosis and treatment. *Deutsches Aerzteblatt International*, 106(34-35), 549-555.
- Teh, C. L., Chew, K. F., & Ling, G. R. (2014). Acute gout in hospitalized patients in Sarawak General hospital. *Medical Journal of Malaysia*, 69(3), 126- 128.
- Terkeltaub, R.A. (2003). Clinical practice gout. *The New England Journal of Medicine*, 349, 1647-1655.
- Torondel, B., Opare, D., Brandberg, B., Cobb, E., & Cairncross, S. (2014). Efficacy of *Moringa oleifera* leaf powder as a hand-washing product: a crossover controlled study among healthy volunteers. *BMC Complementary and Alternative Medicine*, 14(1), 57.
- Triggiani, V., Resta, F., Guastamacchia, E., Sabba, C., Licchelli, B., Ghiyasaldin, S., & Tafaro, E. (2006). Role of antioxidants, essential fatty acids, carnitine, vitamins, phytochemicals and trace elements in the treatment of diabetes mellitus and its chronic complications. *Endocrine, Metabolic & Immune Disorders-Drug Targets*, 6(1), 77-93.
- Truong, D. H., Nguyen, D. H., Ta, N. T. A., Bui, A. V., Do, T. H., & Nguyen, H. C. (2019). Evaluation of the use of different solvents for phytochemical constituents, antioxidants, and *in vitro* anti-inflammatory activities of *Severinia buxifolia*. *Journal of Food Quality*, 2019, 1-9.

- Tsao, R., & Yang, R. (2003). Optimization of a new mobile phase to know the complex and real polyphenolic composition: towards a total phenolic index using high-performance liquid chromatography. *Journal of Chromatography A*, 1018(1), 29-40.
- Tu, S. F., Liu, R. H., Cheng, Y. B., Hsu, Y. M., Du, Y. C., El-Shazly, M., Wu, Y. C., & Chang, F. R. (2014). Chemical constituents and bioactivities of *Clinacanthus nutans* aerial parts. *Molecules*, 19(12), 20382-20390.
- Tuhin, R. H., Begum, M. M., Rahman, M. S., Karim, R., Begum, T., Ahmed, S. U., Mostofa, R., Hossain, A., Abdel-Daim, M., & Begum, R. (2017). Wound healing effect of *Euphorbia hirta* linn. (Euphorbiaceae) in alloxan induced diabetic rats. *BMC Complementary and Alternative Medicine*, 17(1), 423.
- Uboh, F. E., Okon, I. E., & Ekong, M. B. (2010). Effect of aqueous extract of *Psidium guajava* leaves on liver enzymes, histological integrity and hematological indices in rats. *Gastroenterology Research*, 3(1), 32-38.
- Ueda, T., Honda, S., Morikawa, H., Kitamura, S., Iwama, Y., & Nakagawa, K. (2015). Chrysanthemum flower oil inhibits diet-induced serum uric acid elevation in adult male subjects. *Nutrafoods*, 14(3), 151-158.
- Uma, D. B., Ho, C. W., & Wan Aida, W. M. (2010). Optimization of extraction parameters of total phenolic compounds from henna (*Lawsonia inermis*) leaves. *Sains Malaysiana*, 39(1), 119-128.
- Umamaheswari, M., AsokKumar, K., Somasundaram, A., Sivashanmugam, T., Subhadradevi, V., & Ravi, T. K. (2007). Xanthine oxidase inhibitory activity of some Indian medical plants. *Journal of Ethnopharmacology*, 109(3), 547-551.
- Umamaheswari, M., Asokkumar, K., Sivashanmugam, A. T., Remyaraju, A., Subhadradevi, V., & Ravi, T. K. (2009). *In vitro* xanthine oxidase inhibitory activity of the fractions of *Erythrina stricta* Roxb. *Journal of Ethnopharmacology*, 124(3), 646-648.
- Underwood, M. (2006). Diagnosis and management of gout. *BMJ*, 332(7553), 1315-1319.
- Unno, T., Sugimoto, A., & Kakuda, T. (2004). Xanthine oxidase inhibitors from the leaves of *Lagerstroemia speciosa* (L.). *Journal of Ethnopharmacology*, 93(2-3), 391-395.

- Velioglu, Y. S., Mazza, G., Gao, L., & Oomah, B.D. (1998). Antioxidant activity and total phenolics in selected fruits, vegetables, and grain products. *Journal of Agricultural and Food Chemistry*, 46(10), 4113–4117.
- Vitart, V., Rudan, I., Hayward, C., Gray, N. K., Floyd, J., Palmer, C. N., Knott, S.A., Kolcic, I., Polasek, O., Graessler, J., & Wilson, J. F. (2008). SLC2A9 is a newly identified urate transporter influencing serum urate concentration, urate excretion and gout. *Nature Genetics*, 40(4), 437-442.
- Vongsak, B., Sithisarn, P., & Gritsanapan, W. (2013). Simultaneous HPLC quantitative analysis of active compounds in leaves of *Moringa oleifera* Lam. *Journal of Chromatographic Science*, 52(7), 641-645.
- Vongsak, B., Sithisarn, P., Mangmool, S., Thongpraditchote, S., Wongkrajang, Y., & Gritsanapan, W. (2013). Maximizing total phenolics, total flavonoids contents and antioxidant activity of *Moringa oleifera* leaf extract by the appropriate extraction method. *Industrial Crops and Products*, 44, 566-571.
- Wahyuningsih, S., Sukandar, E. Y., & Sukrasno. (2016a). Antihyperuricemia activity of the ethanol extract of Roselle calyx and its fraction (*Hibiscus sabdariffa* Linn) on male wistar rats. *International Journal of Pharmacy and Pharmaceutical Sciences*, 8(3), 278-280.
- Wahyuningsih, S., Sukandar, E. Y., & Sukrasno. (2016b). *In vitro* xanthine oxidase inhibitor activity of ethanol extract and fraction roselle calyx (*Hibiscus sabdariffa* L.). *International Journal of Pharmaceutical and Clinical Research*, 8(6), 619-622.
- Wallace, K. L., Riedel, A. A., Joseph-Ridge, N., & Wortmann, R. (2004). Increasing prevalence of gout and hyperuricemia over 10 years among older adults in a managed care population. *The Journal of rheumatology*, 31(8), 1582-1587.
- Wan, R. W., Mahfudzah, A., Nazihah, M. Y., Tan, H. L., Wan, S. W., Amanda, J. P., & Tony, R. M. (2018). Association of solute carrier family 2, member 9 (SLC2A9) genetic variant rs3733591 with gout in a Malay sample set. *The Medical Journal of Malaysia*, 73(5), 307-310.
- Wang, Y., Zhu, J. X., Kong, L. D., Yang, C., Cheng, C. H. K., & Zhang, X. (2004). Administration of procyanidins from grape seeds reduces serum uric acid levels and decreases hepatic xanthine dehydrogenase/oxidase activities in oxonate-treated mice. *Basic & Clinical Pharmacology & Toxicology*, 94(5), 232-237.



- Wang, Y., Deng, T., Lin, L., Pan, Y., & Zheng, X. (2006). Bioassay-guided isolation of antiatherosclerotic phytochemicals from *Artocarpus altilis*. *Phytotherapy Research: An International Journal Devoted to Pharmacological and Toxicological Evaluation of Natural Product Derivatives*, 20(12), 1052-1055.
- Wang, J., Sun, B., Cao, Y., Tian, Y., & Li, X. (2008). Optimisation of ultrasound-assisted extraction of phenolic compounds from wheat bran. *Food Chemistry*, 106(2), 804-810.
- Wang, M., Cao, A., Ouyang, C., Li, Y., & Wei, Y. (2015). Rapid screening and identification of non-target flavonoid components in invasive weeds by LC/MS-IT-TOF. *Analytical Methods*, 7(24), 10207-10216.
- Widyarini, K. D., Sukandar, E. Y., & Fidrianny, I. (2015). Xanthine oxidase inhibitory and antihyperuricemic activities of *Anredera cordifolia* (Ten) Steenis, *Sonchus arvensis* L. and its combination. *International Journal of Pharmacy and Pharmaceutical Sciences*, 7(3), 87-90.
- Wikimedia commons. (2014). *Euphorbia hirta*. Retrieved March 19, 2019 from [https://commons.wikimedia.org/wiki/File:Asthma\\_Weed\\_\(Euphorbia\\_hirta\)\\_4.jpg](https://commons.wikimedia.org/wiki/File:Asthma_Weed_(Euphorbia_hirta)_4.jpg)
- Wikimedia commons. (2017). *Artocarpus altilis*. Retrieved May 11, 2020 from [https://commons.wikimedia.org/wiki/File:Artocarpus\\_altilis\\_\(Moraceae\).jpg](https://commons.wikimedia.org/wiki/File:Artocarpus_altilis_(Moraceae).jpg)
- Wong, Y. P., Ng, R. C., Chuah, S. P., Koh, R. Y., & Ling, A. P. K. (2014). Antioxidant and xanthine oxidase inhibitory activities of *Swietenia macrophylla* and *Punica granatum*. *International Conference on Biological, Environment and Food Engineering*. Bali, Indonesia. 53-58.
- Wu, X. W., Lee, C. C., Muzny, D. M., & Caskey, C. T. (1989). Urate oxidase: primary structure and evolutionary implications. *Proceedings of the National Academy of Sciences*, 86(23), 9412-9416.
- Wu, L., Li, Y. L., Li, S. M., Yang, X. W., Xia, J. H., Zhou, L., & Zhang, W. D. (2010). Systematic phytochemical investigation of *Abies spectabilis*. *Chemical and Pharmaceutical Bulletin*, 58(12), 1646-1649.
- Wu, X. Y., Tang, A. C., & Lu, Q. Y. (2012). Study on antitumor effect of the extract from *Christia vespertilionis* in vivo. *Chinese Journal of Experimental Traditional Medical Formulae*, 8, 202-204.

- Yaacob, N. S., Hamzah, N., Kamal, N. N. N. M., Abidin, S. A. Z., Lai, C. S., Navaratnam, V., & Norazmi, M. N. (2010). Anticancer activity of a sub-fraction of dichloromethane extract of *Strobilanthes crispus* on human breast and prostate cancer cells *in vitro*. *BMC Complementary and Alternative Medicine*, 10(1), 42.
- Yan, H., Ma, Y., Liu, M., & Zhou, L. (2008). The dual actions of *Paederia scandens* extract as a hypouricemic agent: xanthine oxidase inhibitory activity and uricosuric effect. *Planta Medica*, 74(11), 1345-1350.
- Yan, J., Zhang, G., Hu, Y., & Ma, Y. (2013). Effect of luteolin on xanthine oxidase: Inhibition kinetics and interaction mechanism merging with docking simulation. *Food Chemistry*, 141(4), 3766-3773.
- Yan, M., An, Y. T., Li, J., Wu, Z. Z., & Wang, T. (2014). Regulatory effect of leonurus extracts on hyperuricemia in rats. *China Journal of Chinese Materia Medica*, 39(24), 4856-4859.
- Yazid, E. A., Supriyanto, G., & Tjahjandarie, T. S. (2014). Development of spectrophotometric method for allopurinol in urine based on the diazotization reaction. *Malaysian Journal of Analytical Sciences*, 18(1), 212-220.
- Yeap, S. S., Goh, E. M., & Gun, S. C. (2009). A survey on the management of gout in Malaysia. *International Journal of Rheumatic Diseases*, 12(4), 329-335.
- Yerlikaya, A., Dagel, T., King, C., Kuwabara, M., Lanaspa, M. A., Andres-Hernando, A., Covic, A., Manitius, J., Sag, A. A., & Kanbay, M. (2017). Dietary and commercialized fructose: Sweet or sour?. *International Urology and Nephrology*, 49(9), 1611-1620.
- Yilmaz, Y., & Toledo, R. T. (2006). Oxygen radical absorbance capacities of grape/wine industry byproducts and effect of solvent type on extraction of grape seed polyphenols. *Journal of Food Composition and Analysis*, 19(1), 41-48.
- Yoon, I. S., Park, D. H., Bae, M. S., Oh, D. S., Kwon, N. H., Kim, J. E., Choi, C. Y., & Cho, S. S. (2017). *In vitro* and *in vivo* studies on *Quercus acuta* Thunb.(Fagaceae) extract: Active constituents, serum uric acid suppression, and xanthine oxidase inhibitory activity. *Evidence-based Complementary and Alternative Medicine*, 2017, 1-9.

- Yumita, A., Suganda, A. G., & Sukandar, E. Y. (2013). Xanthine oxidase inhibitory activity of some Indonesian medicinal plants and active fraction of selected plants. *International Journal of Pharmacy and Pharmaceutical Sciences*, 5, 293-296.
- Zakaria, M., & Mohd, M.A. (2010). *Traditional Malay medicinal plants*. Kuala Lumpur: Institut Terjemahan Negara Malaysia.
- Zakaria, F., Ibrahim, W. N. W., Ismail, I. S., Ahmad, H., Manshoor, N., Ismail, N., Zainal, Z., & Shaari, K. (2019). LCMS/MS metabolite profiling and analysis of acute toxicity effect of the ethanolic extract of *Centella asiatica* on zebrafish model. *Pertanika Journal of Science & Technology*, 27(2), 985-1003.
- Zain, A.A., & Suresh, S. (2015). *Arthritis in Men*. Retrieved on October 13, 2016, from <https://www.bfm.my/podcast/the-bigger-picture/health-and-living/amir-azlan-zain-suhail-suresh-gout-uric-acid>
- Zehl, M., Braunberger, C., Conrad, J., Crnogorac, M., Krasteva, S., Vogler, B., Beifuss, U., & Krenn, L. (2011). Identification and quantification of flavonoids and ellagic acid derivatives in therapeutically important *Drosera* species by LC–DAD, LC–NMR, NMR, and LC–MS. *Analytical and Bioanalytical Chemistry*, 400(8), 2565-2576.
- Zeng, Y. C., Li, S., Liu, C., Gong, T., Sun, X., Fu, Y., & Zhang, Z.R. (2017). Soluplus micelles for improving the oral bioavailability of scopoletin and their hypouricemic effect *in vivo*. *Acta Pharmacologica Sinica*, 38(3), 424-433.
- Zhang, Y., Chen, C., Choi, H., Chaisson, C., Hunter, D., Niu, J., & Neogi, T. (2012). Purine-rich foods intake and recurrent gout attacks. *Annals of the Rheumatic Diseases*, 71(9), 1435-1436.
- Zhang, J., Shuai, X., Li, J., Xiang, N., Gong, T., & Zhang, Z. (2016a). Biodistribution, hypouricemic efficacy and therapeutic mechanism of morin phospholipid complex loaded self-nanoemulsifying drug delivery systems in an experimental hyperuricemic model in rats. *Journal of Pharmacy and Pharmacology*, 68(1), 14-25.
- Zhang, H. J., Hu, Y. J., Xu, P., Liang, W. Q., Zhou, J., Liu, P. G., Cheng, L., & Pu, J. B. (2016b). Screening of potential xanthine oxidase inhibitors in *Gnaphalium hypoleucum* DC. by immobilized metal affinity chromatography and ultrafiltration-ultra performance liquid chromatography- mass spectrometry. *Molecules*, 21(9), 1242.

- Zhao, X., Zhu, J. X., Mo, S. F., Pan, Y., & Kong, L. D. (2006). Effects of cassia oil on serum and hepatic uric acid levels in oxonate-induced mice and xanthine dehydrogenase and xanthine oxidase activities in mouse liver. *Journal of Ethnopharmacology*, 103(3), 357-365.
- Zhao, F., Guochun, L., Yang, Y., Shi, L., Xu, L., & Yin, L. (2015). A network pharmacology approach to determine active ingredients and rationality of herb combinations of Modified-Simiaowan for treatment of gout. *Journal of Ethnopharmacology*, 168, 1-16.
- Zhishen, J., Mengcheng, T., & Jianming, W. (1998). The determination of flavonoid contents in mulberry and their scavenging effects on superoxide radicals. *Food Chemistry*, 64, 555-559.
- Zhou, K., & Yu, L. (2006). Total phenolic contents and antioxidant properties of commonly consumed vegetables grown in Colorado. *Food Science and Technology*, 39(10), 1155-1162.



PTTA UTHM  
PERPUSTAKAAN TUNKU TUN AMINAH